Fwd: [Cag-all-users] Call for articles for the Second Issue of Journal of the Indian Audit & Accounts Department (IA&AD)- Manthan – the Journal of Public Sector Auditing and Accounting of SAI India

RAJESH KUMAR NARWAL Sr Audit Officer Civil < narwalrk.hmp.sca@cag.gov.in >

Thu, 01 Jan 2026 5:20:58 PM +0530

To "Himanshu Juneja"<himanshuj.guj.au@cag.gov.in>,"Abhinesh Meena"<abhineshm.hmp.au@cag.gov.in>

Please upload on official website

========== Forwarded message ==========
From: NAAA Shimla via Cag-all-users <cag-all-users@lsmgr.nic.in>
To: "cag-all-users"<cag-all-users@lsmgr.nic.in>, "CAG-ALL-OFFICES"<cag-all-offices@lsmgr.nic.in>, "cag-iaas"<cag-iaas@lsmgr.nic.in>
Date: Thu, 01 Jan 2026 16:18:56 +0530
Subject: [Cag-all-users] Call for articles for the Second Issue of Journal of the Indian Audit & Accounts Department (IA&AD)- Manthan – the Journal of Public Sector Auditing and Accounting of SAI India
========== Forwarded message ==========

Madam/Sir,

New Year Greetings from the National Academy of Audit and Accounts (NAAA), Shimla!

Following the successful launch of the first issue of *Manthan – The Journal of Public Sector Auditing and Accounting of SAI India* by the Comptroller and Auditor General of India on 28 November 2025, we are pleased to inform you that NAAA is now in the process of bringing out the **second issue** of the journal. We invite contributions from authors across the IA&AD for the upcoming edition.

Articles from experts in universities and professional institutes (IITs, IIMs, ICAI, ICMAI, etc.) are also welcome, provided they arise out of joint research or collaborative projects related to the IA&AD's mandate. This will help ensure that the journal's core focus on public sector auditing and accounting is maintained.

We have prepared an indicative list of sectors on which the articles may focus, as well as detailed guidelines for writing and submitting manuscripts. These documents are attached for your reference. We look forward to your contributions in this collective effort to promote knowledge exchange within the IA&AD and beyond.

We would be deeply grateful for your participation in this collaborative academic initiative. Your experience, analytical insight, and professional expertise hold immense value for the wider auditing and accounting community. Each contribution has the potential to inspire, inform and guide both practitioners and scholars. **We would be honoured to include your perspectives in this edition and believe that your participation will greatly enrich the journal's intellectual depth.**

To streamline and facilitate the submission process, we have enabled online submission. Kindly follow the steps below:

1. Register as an Author
Please fill in the required details here to register as an author:
http://10.47.37.26/ojs/index.php/Manthan/user/register

Please note that the link usually works only within the IA&AD network. If you are unable to access it, kindly send your article and annexures, by email to ***naaa@cag.gov.in****, with a copy to* ***dasnd@cag.gov.in***

2. Author Undertaking Form
Manthan is a peer-reviewed journal and adheres to standard ethical guidelines. Please fill out, sign and return the attached undertaking form confirming the originality of your work and compliance with ethical protocols.

We request you to submit your article along with the completed undertaking form by 15 January 2026. A sample article is also attached for your guidance on formatting and content structure.

Should you have any queries, please feel free to reach out to us at **naaa@cag.gov.in** / **naaa.cag@nic.in**, with a copy to **dasnd@cag.gov.in**

भवदीय,
नन्द दुलाल दास
निदेशक (प्रशिक्षण-I)
राष्ट्रीय लेखापरीक्षा एवं लेखा अकादमी, शिमला

========== Forwarded message ==========
From: NAAA Shimla <naaa@cag.gov.in>
To: "cag-all-users"<cag-all-users@lsmgr.nic.in>, "CAG-ALL-OFFICES"<cag-all-offices@lsmgr.nic.in>, "cag-iaas"<cag-iaas@lsmgr.nic.in>
Date: Sat, 06 Dec 2025 16:28:17 +0530
Subject: Call for articles for the Second Issue of Journal of the Indian Audit & Accounts Department (IA&AD)- Manthan – the Journal of Public Sector Auditing and Accounting of SAI India
========== Forwarded message ==========

Madam/Sir,

Greetings from the National Academy of Audit and Accounts (NAAA), Shimla!

Following the successful launch of the first issue of *Manthan – The Journal of Public Sector Auditing and Accounting of SAI India* by the Comptroller and Auditor General of India on 28 November 2025, we are pleased to inform you that NAAA is now in the process of bringing out the **second issue** of the journal. We invite contributions from authors across the IA&AD for the upcoming edition.

Articles from experts in universities and professional institutes (IITs, IIMs, ICAI, ICMAI, etc.) are also welcome, provided they arise out of joint research or collaborative projects

Guidelines for the Authors for Registration:

- Fill out the Registration form at: http://10.47.37.26/ojs/index.php/Manthan/user/register
- **Registration Guidelines:** Kindly register as an author with all the necessary details in the link provided for registration. **Do not check on the input that mentions '**I would like to be contacted with requests to review submissions to this journal.'



Please note that the link usually works only within the IA&AD network. If you are unable to access it, kindly send your article and annexures, by email to ***naaa@cag.gov.in****, with a copy to* ***dasnd@cag.gov.in***

Article Writing Guidelines:

- Clear writing depends on clear thinking. Please think and prepare an outline about the article before writing the article.
- The article should be written in such a way that it represents an analysis done by the author and not mere reproduction of facts and figures.
- All the facts, figures and tables should be appropriately cited, unless created by the author.
- APA style of referencing may be followed in all articles. Please see how the citation reads.
- The size of the article may be restricted to a maximum of 5 pages, with font size 12 (Times New Roman and multiple 1.5 line spacing; paragraph spacing: 6 points before and after).
- Please complete the article within given timeline and send to the journal for review to the NAAA/editors/ their nominee by email.
- After receipt of the comments of the peer-reviewer, the revisions may be carried out within two weeks and resent to the NAAA/editors/ their nominee by email.

Author's Undertaking Form

Journal Title: Manthan: Journal of Public Sector Auditing and Accounting of SAI India

Article Title: [Insert Full Title of Article]

Corresponding Author: [Insert Full Name of Corresponding Author]

This form must be completed and signed by all authors of the submitted manuscript. By signing this form, each author acknowledges and agrees to the following:

1. Originality and Authorship Confirmation

- I/We confirm that this manuscript is original work and has not been published elsewhere in any language, nor is it under consideration for publication elsewhere.
- I/We confirm that any individual who contributed to the work but does not meet the criteria for authorship (e.g., technical assistance, writing assistance, general support) has been acknowledged, with their permission, in the "Acknowledgements" section.

2. Ethical Conduct and Research Integrity

- I/We confirm that the research presented in this manuscript was conducted in an ethical and responsible manner.
- For primary data collection/ studies involving human participants, I/we confirm that appropriate institutional review board (IRB) or equivalent ethics committee approval was obtained, and that informed consent was secured from all participants (or their legal guardians). The manuscript includes a statement to this effect.
- For studies involving animals, I/we confirm that all institutional and national guidelines for the care and use of laboratory animals were followed. The manuscript includes a statement to this effect.
- I/We confirm that the data presented in this manuscript are authentic and accurate, and that no data fabrication, falsification, or manipulation has occurred.
- I/We understand that plagiarism, self-plagiarism, redundant publication, and other forms of research misconduct are strictly prohibited. I/We confirm that this manuscript is free from such misconduct.

3. Conflict of Interest Disclosure

- I/We declare any actual or potential financial or non-financial conflicts of interest that could be perceived to influence the results or interpretation of this manuscript.
- If no conflicts of interest exist, I/we explicitly state "The authors declare no conflict of interest." in the manuscript.
- If conflicts of interest exist, I/we have provided a detailed statement in the manuscript under a "Conflicts of Interest" section, fully disclosing the nature of the conflict.

4. Funding Disclosure

- I/We acknowledge all sources of financial support for the work, including grants, scholarships, and institutional support. These sources have been fully disclosed in an "Funding" section within the manuscript.

5. Copyright and Permissions

- I/We warrant that the manuscript does not infringe upon any copyright or other proprietary rights of any third party.
- I/We have obtained written permission from the copyright holder for any previously published material (e.g., figures, tables, extensive text excerpts) included in this manuscript that is not in the public domain or covered by fair use/dealing. Copies of these permissions will be provided to the journal upon request.
- I/We understand that upon acceptance, copyright may be transferred to the journal/publisher, or the article may be published under an open access license (e.g., Creative Commons), as per the journal's policy.

6. Data Availability and Transparency

- I/We agree to make the raw data, methods, and materials underlying the findings described in this manuscript available to the journal and, where appropriate and ethical, to the scientific community for verification purposes upon reasonable request.
- I/We confirm that any data sharing adheres to principles of privacy and confidentiality, especially for studies involving human subjects.

8. Undertaking and Declaration I/We, the undersigned authors, have read and understood the terms of this Author Undertaking Form and agree to abide by them. I/We understand that any breach of these declarations may lead to the withdrawal or retraction of the manuscript or published article.

Author Signatures:

Sr. No.	Full Name	Signature	Date	Email
1				
2				
3				

Sr. No.	Sector Name /Area	
1	Agriculture and Allied Sectors	
2	Mining	1. Surface Mining 2. Underground Mining
3	Education:	1. Elementary 2. Secondary 3. Higher
4	Local Bodies	1. Urban Local Bodies 2. Rural Local Bodies
5	Transport	1. Road 2. Rail 3. Air 4. Water
6	Service Sector	1. IT Service 2. IT Security 3. IT Audit 4. AI and ML 5. GIS & Remote Sensing 6. Logistic and Infrastructure 7. Health 8. Telecommunication
7	Trade	1. National 2. International
8	Energy	1. Renewable 2. Non-Renewable
9	Tourism Sector	1. National 2. International 3. Eco Tourism
10	Industries	1. Heavy 2. MSME
11	Social Benefits	1. Subsidies 2. Public Distribution System 3. Social Upliftment schemes
12	Environmental Audit	
13	Gross National Happiness (GNH)	
14	National Income and Indices Accounting	
15	Child Labour and Bonded Labour	
16	UN and Satellite Organisations	
17	Public Sector	
18	Company Accounts	
19	Government Accounts	
20	Entitlements: Pension and GPF	

Marine and Coastal Fisheries, 2025, **17**, vtaf009
https://doi.org/10.1093/mcfafs/vtaf009
Advance access publication: May 27, 2025
Review

Science priorities to evaluate the effects of offshore wind energy development on fish and fisheries in the Gulf of America

Shayna A. Sura[1,*], Raymond E. Czaja Jr.[1], Nathan Brugnone[2,3,**], Sarah L. Gibbs[4,5,**], J. Read Hendon[6,**], Willem Klajbor[7,8,9,**], Avery B. Paxton[10,**], R. Ryan Rindone[11,**], Skyler R. Sagarese[10,**], Kate Wing[12,**], Leann Bosarge[13,**], David D. Chagaris[14,**], William D. Heyman[15,**], Michelle A. Johnston[16,**], James A. Morris Jr.[17,**], William F. Patterson III[18,**], James M. Tolan[**,‡], John F. Walter III[9,**], and Holden E. Harris[7,9]

[1]National Center for Ecological Analysis and Synthesis, University of California Santa Barbara, Santa Barbara, California, USA
[2]Two Six Technologies, Arlington, Virginia, USA
[3]Michigan State University, East Lansing, Michigan, USA
[4]Stokes School of Marine and Environmental Sciences, University of South Alabama, Mobile, Alabama, USA
[5]Dauphin Island Sea Lab, Dauphin Island, Alabama, USA
[6]National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southeast Fisheries Science Center, Pascagoula, Mississippi, USA
[7]Cooperative Institute of Marine and Atmospheric Studies, Rosenstiel School of Marine Atmospheric and Earth Science, University of Miami, Miami, Florida, USA
[8]National Oceanic and Atmospheric Administration, Atlantic Oceanographic and Meteorological Laboratory, Miami, Florida, USA
[9]National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southeast Fisheries Science Center, Miami, Florida, USA
[10]National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southeast Fisheries Science Center, Beaufort, North Carolina, USA
[11]Gulf of Mexico Fishery Management Council, Tampa, Florida, USA
[12]Intertidal Agency, Berkeley, California, USA
[13]Bosarge Boats, Inc., Pascagoula, Mississippi, USA
[14]Nature Coast Biological Station, Institute of Food and Agriculture Sciences, University of Florida, Gainesville, Florida, USA
[15]LGL Ecological Research Associates, Inc., Bryan, Texas, USA
[16]National Oceanic and Atmospheric Administration, Flower Garden Banks National Marine Sanctuary, Galveston, Texas, USA
[17]National Oceanic and Atmospheric Administration, National Ocean Service, National Centers for Coastal Ocean Science, Beaufort, North Carolina, USA
[18]School of Forest, Fisheries, and Geomatic Sciences, University of Florida, Gainesville, Florida, USA

*Corresponding author: Shayna A. Sura. Email: ssura@ucsb.edu.
**These authors contributed equally and are ordered alphabetically
‡Retired.

ABSTRACT

Objective: Offshore wind (OSW) farms are slated for development in the Gulf of America (also known as the Gulf of Mexico), presenting a timely need to understand the potential effects of their construction and operation on marine ecosystems.

Methods: To help address this need, we convened a transdisciplinary working group of scientists, managers, and representatives of commercial and recreational fisheries to identify and assess research priorities and recommendations related to the effects of OSW farms on fish and fisheries in the Gulf of America.

Results: Here, we share these research priorities for shrimp, reef fishes, coastal migratory pelagics, forage fishes, oceanic pelagic fishes, coastal elasmobranchs, and invasive species. We then detail OSW research needs that are related to oceanographic and ecological processes, and we provide specific recommendations for fisheries management, marine spatial planning, and detection of social and economic effects. Our synthesis highlights three overarching considerations: (1) targeted data collection is needed to disentangle the effects of OSW from those of concurrent natural and anthropogenic stressors, (2) measuring the effects of OSW will require maintaining the integrity of

Received: November 2, 2024. **Revised:** January 30, 2025. **Editorial decision:** February 8, 2025
© The Author(s) 2025. Published by Oxford University Press on behalf of American Fisheries Society.
This is an Open Access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

long-term fisheries-independent surveys and augmenting such surveys with comprehensive before-after-control-impact or before-after-gradient research designs, and (3) there are differences in public participation processes for nascent OSW development versus established fisheries management that should be considered to allow meaningful societal participation.

Conclusions: Scientists and natural resource managers have a unique opportunity to address these priorities and recommendations, shaping the understanding of the effects of OSW.

KEYWORDS: artificial reefs, fisheries, fisheries management, Gulf of America, marine artificial structures, offshore wind, research priorities

LAY SUMMARY

Given the multisector (oil and gas, artificial reefs, fisheries) and multispecies fisheries nature of the United States Gulf of America, we summarize important research priorities for understanding the effects of offshore wind development on fish and fisheries.

INTRODUCTION

The rapid and global expansion of offshore wind (OSW) development presents pressing science and management opportunities and questions. Since the first OSW farm (Vindeby) was built in 1991 in the North Sea (Ørsted, 2019), OSW farms have been developed in northwest Europe (Ørsted, 2019; Wang et al., 2024), China (Wang et al., 2024), and the United States (Doty, 2023). Europe led the global development of OSW until 2022, when the Asia-Pacific region began producing over 34 gigawatts (GW) per year from OSW (Wang et al., 2024; Williams et al., 2023). As of 2022, U.S. OSW contributes to <1% of global OSW electricity generation (Williams et al., 2023). Over the past decade, the USA has set federal and regional plans to increase technological capacity, reduce production costs, and expand OSW infrastructure (The White House, 2021; Musial et al., 2023; U.S. Department of Energy, 2022); however, these plans have more recently been curtailed (The White House, 2025).

The Gulf of America (hereafter, abbreviated as GoA; also known as Gulf of Mexico) has been explored as a suitable region for development of OSW in the USA (Randall et al., 2022). The Bureau of Ocean Energy Management (BOEM), under the U.S. Department of the Interior, executed the first federal offshore wind energy lease sale in the GoA in 2023 and has prepared other wind energy areas (WEAs) in U.S. federal waters off Texas and Louisiana for potential auction in the future (Figure 1). Other GoA areas that are under consideration and review for OSW leasing have been identified (Musial et al., 2023), and two additional leases in Louisiana state waters have been designated and awarded (Baurick, 2023). Given an estimated timeline of approximately 8 years from leases being granted to the beginning of infrastructure installation (Figure 2; Bureau of Ocean Energy Management, 2024), the GoA may have operational OSW infrastructure in the early 2030s. Therefore, the time to anticipate the potential ecological impacts of OSW in the GoA, especially regarding fish and fisheries, and plan for their mitigation is now.

Offshore wind infrastructure in the U.S. GoA would join a constellation of oil and gas (O&G) platforms, pipelines, and intentionally placed artificial reefs in a heavily industrialized and immensely productive ecosystem. Collectively, the GoA has the largest footprint of artificial reefs in the USA (Paxton et al., 2024) and contains 49% of the global area of offshore O&G platforms (Bugnot et al., 2021). Thousands of O&G platforms have been constructed since the 20th century (Shipp & Bortone, 2009; Yergin, 1991), with about 2,100 platforms existing currently (Office for Coastal Management, 2024). Also, over 20,000 documented (and innumerable undocumented) artificial structures have been deployed on the GoA shelf (Gardner et al., 2022; Schulze et al., 2020; Shipp & Bortone, 2009). In addition, the GoA provides 17% of the U.S. national commercial fisheries landings, 35% of the U.S. national recreational fisheries catch, and 21% of the U.S. marine aquaculture production value (National Marine Fisheries Service [NMFS], 2024a). Artificial structures support commercial and recreational GoA fisheries (Gallaway et al., 2009, 2021), and fishermen in the GoA have traditionally targeted fishing on O&G platforms and artificial reefs (Brashier, 1988; Stanley & Wilson, 1989). The presence of multisector and multispecies fisheries in the GoA presents novel issues, research opportunities, and management scenarios to consider with respect to the potential effects of OSW on fish and fisheries.

Approach

We convened a transdisciplinary working group in 2023 (Impacts of Offshore Wind on Gulf of Mexico Fish and Fisheries, https://www.nceas.ucsb.edu/workinggroups/gei-data-synthesis-and-models-evaluate-cumulative-ecosystem-impacts-offshore-wind) to evaluate potential effects of OSW development on fish and fisheries in the GoA. The working group members represented academic, federal, and private research entities; federal and state natural resource management agencies; and recreational and commercial fisheries groups (Table 1). We used a coproduction process (e.g., Beier et al., 2017) to identify research priorities and rank them according to urgency and importance (Figure 3). Although our primary focus was on fish and fisheries, we also acknowledge the importance of identifying and addressing research priorities for other ecologically important and/or protected species (e.g., birds, sea turtles, and marine mammals). In addition, OSW in the GoA may become coupled with "green hydrogen" technology in the future (Hicks, 2024), as the GoA is one region where low costs of production could be achieved (Brunik et al., 2024); however, considering the effects of green hydrogen technology was outside the scope of our working group.

Here, we present the priorities and recommendations for GoA managers that were identified by the working group (Table 2). In section 2, we present research priorities for specific taxa and fisheries of interest in the GoA. In section 3, we more broadly discuss research priorities for oceanographic and

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025



Figure 1. Current and proposed OSW development in the GoA for (A) federal waters and (B–C) state waters. (A) Federal offshore wind development activities, including the regional wind energy call area (gray outline), potential wind energy areas (teal polygons), proposed lease areas (pink outlines), and current wind lease (yellow polygon). Potential wind energy areas (teal polygons) denote options for wind energy development created by the Bureau of Ocean Energy Management and the National Oceanic and Atmospheric Administration's National Centers for Coastal Ocean Science. The current wind lease (yellow polygon) was awarded to RWE (Essen, Germany) in October 2023. The unsolicited lease requests (pink outlines) are from Hecate Energy (Chicago, Illinois), announced in July 2024. Extent indicators correspond to state leases (purple outline; orange outline) for (B) Diamond Offshore Wind (Boston, Massachusetts; purple polygon) and for (C) Vestas Offshore Wind (Aarhus, Denmark; orange polygon). Bathymetry map (inset) is from the General Bathymetric Chart of the Oceans (GEBCO); the National Oceanic and Atmospheric Administration's National Centers for Environmental Information. The locations of other artificial structures, including (D) artificial reefs and (E) oil and gas platforms, are also provided. Artificial reef and oil and gas platform data were downloaded from Marine Cadastre (hub.marinecadastre.gov) on July 19, 2024.



Figure 2. Timeline of offshore wind infrastructure development in the Gulf of America. The major phases include planning and analysis, leasing, site assessment, construction, operations, and decommissioning. The numbers indicate an estimated time frame in years for each phase. The image was provided by the Bureau of Ocean Energy Management (BOEM). Abbreviations in the image are as follows: SAP = site assessment plan; COP = construction and operations; BSEE = Bureau of Safety and Environmental Enforcement; FDR/FIR = facility design report/fabrication and installation report.

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Table 1. Representation across scientific, management, and resource/fisheries sectors for members of the Impacts of Offshore Wind on Gulf of Mexico Fish and Fisheries Working Group.

Sector	Working group member affiliations
Academic research	• University of Florida • University of Miami Cooperative Institute of Marine and Atmospheric Sciences • National Center for Ecological Analysis and Synthesis • University of South Alabama
Federal research	• National Oceanic and Atmospheric Administration (NOAA), National Marine Fisheries Service • Bureau of Ocean and Energy Management
Private research	• LGL Ecological Research Associates • TwoSix Technologies • Intertidal Agency
Federal and state natural resource management	• Gulf of Mexico Fishery Management Council (GMFMC) • Texas Parks and Wildlife Coastal Fisheries Division • Louisiana Department of Wildlife and Fisheries • NOAA Office of National Marine Sanctuaries
Recreational and commercial fisheries	• Southern Shrimp Alliance • GMFMC as recounting interests of recreational and commercial fishers • Working group members as individual recreational anglers

ecological processes. In section 4, we discuss research priorities for understanding the social and management implications of OSW development. Finally, we highlight the importance of comprehensive assessment and management processes, as OSW development coincides with climate change and other existing stressors in this large marine ecosystem. The research priorities and recommendations that are presented here will help to ensure an evidence-based approach to understanding and evaluating potential effects of OSW on fish and fisheries in the GoA.

GULF OF AMERICA TAXA AND FISHERIES OF CONCERN WITH RESPECT TO OSW DEVELOPMENT

Myriad coastal and marine habitats in the GoA host an estimated 1,443 species of fish from 700 genera and 223 families (McEachran & Fechhelm, 2006). Many of these species associate with artificial structures, including O&G platforms, or otherwise occur in continental shelf waters that host energy extraction platforms. Approximately 50 finfish species and an additional 11 shellfish species support notable commercial or recreational fisheries (Ward & Tunnell, 2017). In this section, we identify research questions for taxonomic groups or fisheries as they relate to potential effects (and pathways for ecological and social impacts) of OSW (Table 3).

Penaeid shrimp (Table 3)

Commercial bottom trawling of penaeid shrimp—brown shrimp *Farfantepenaeus aztecus*, pink shrimp *F. duorarum*, and white shrimp *Litopenaeus setiferus*—collectively comprise the most valuable commercial fishery in the GoA, with more than $US300 million in dockside value in 2022 (National Marine



Figure 3. Eisenhower diagram with a horizontal axis of urgency from high to low (left to right) and a vertical axis of importance from high to low (top to bottom). Credit: Impacts of Offshore Wind on Gulf of Mexico Fish and Fisheries Working Group attendees, unpublished.

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Table 2. Table of contents for remaining sections of this article. OSW = offshore wind.

Section	Topics
Section 2. Gulf of America taxa and fisheries of concern with respect to OSW development	• Penaeid shrimp • Reef fishes • Coastal migratory pelagic fishes • Forage fishes • Oceanic pelagic fishes • Coastal elasmobranchs • Invasive species

Section 3. Overarching oceanographic and ecological processes	• Oceanographic processes
	  • Wind
	  • Hypoxia and stratification
	  • Benthic and biogeochemical processes
	• Ecological processes
	  • Addition of new artificial hard-bottom and vertical habitat
	  • Succession
	  • Habitat
	  • Connectivity

Section 4. Management and human dimensions implications	• Fisheries management • Marine spatial planning • Social and economic impacts
Section 5. Discussion	• Disentangling OSW from concurrent environmental pressures • Importance of baseline data • Societal participation in the scientific process

Fisheries Service Office of Science and Technology). Catches of all three species occur on the GoA continental shelf, with brown shrimp catches being largest in federal waters off Texas (Tunnell, 2017; Williams et al., 2024) that coincide with the first GoA federal OSW wind energy areas. In these fishing grounds, OSW farms will create de facto spatial closures to the commercial shrimp fishery because trawling vessels may be unable to physically navigate trawlable bottom within the footprint of wind farms. Members of the shrimp fishery have expressed interest in collaborative efforts with researchers to collect in situ data for assessing the ecological impacts of OSW on GoA fisheries. Currently, the precise amount of trawling effort that would be displaced by OSW, the effects on income, and the collaborative research opportunities remain unknown. Future distributions of fisheries compensation funds would also benefit from a regional fund administrator to streamline and standardize the application and approval processes, similar to an effort that is currently underway in the northeast USA by the New York State Energy Research and Development Authority (South Atlantic Fishery Management Council, 2024; Special Initiative on Offshore Wind, 2024).

Penaeid shrimp species in the GoA feed on detritus, algae, and invertebrates, including polychaetes and amphipods (Cook & Lindner, 1970; Pattillo et al., 1997). The availability of these food items may increase near OSW structures due to enhanced biological production from colonizing sessile epifauna, providing a potential positive ecological effect of OSW on benthic macrofauna, including shrimp (Lefaible et al., 2023; Raoux et al., 2017). In the North Atlantic, Krone et al. (2017) estimated that OSW structures may increase the production of commercially important decapods (e.g., crabs *Cancer* spp.), although the role of food availability is unknown. Other researchers found that lobster *Homarus* spp. abundances do not change near OSW structures (Roach et al., 2022; Wilber et al., 2024). Differences in habitat requirements, among other characteristics (e.g., anatomical, physiological, and behavioral traits), of decapods may drive species-specific responses to OSW structures.

There is also potential for OSW development to cause negative habitat alterations for penaeids, including altered sediment composition and direct physical disturbances. Penaeids generally prefer open, soft sediment bottom habitat with relatively little hard structure present (Williams, 1958). Although Hutchison et al. (2020b) found no strong changes in sediment grain size within 90 m of OSW turbines in the northwest Atlantic, potential changes in sediment composition in the GoA cannot be ruled out. The negative effects of direct physical disturbances from OSW, such as electric and magnetic fields (EMF) and noise, on decapod behavior and physiology are relatively well documented (Harsanyi et al., 2022; Leiva et al., 2021; Scott et al., 2018). However, no investigations have tested such adverse outcomes for penaeid shrimp. Also, GoA penaeid shrimp exhibit species-specific traits and behaviors; for example, the burrowing behavior of brown shrimp differs from that of white shrimp (Osborn et al., 1969). This emphasizes the need for species-specific research to understand the effects of OSW-related physical disturbances on GoA penaeid shrimp. Additionally, there is concern that thermal emissions from cables (among other OSW activities) may affect shrimp, as penaeid growth, abundance, and distribution are temperature dependent (Arreguín-Sánchez et al., 2015; Montero et al., 2016; Zein-Eldin & Aldrich, 1965). Currently, it is unknown whether OSW cables produce localized temperature alterations; thus, benthic environmental monitoring may clarify this potential issue, among others.

Many study types are needed to evaluate shrimp responses to OSW development. Shrimp survival, growth, and behavior are largely tied to benthic environmental conditions (Haas et al., 2004; Zein-Eldin & Renaud, 1986); thus, monitoring before, during, and after operational phases of OSW development would aid in detecting changes. Collecting baseline benthic environmental conditions, including the presence of EMF and substrate temperature and condition, before OSW development would ideally include control sites to discriminate widespread change from changes caused by OSW development. Observational field studies may detect population-level changes (e.g., shrimp abundance) in response to potential changes in the benthic environment. Manipulative lab experiments may detect individual-level effects of physical disturbances (e.g., effects of noise on burrowing behavior in shrimp), which could identify potential mechanisms for observed population-level trends. Hybrid field experiments that quantify direct disturbances (e.g., EMF and heat from cables) and use cages to manipulate in situ crustacean exposure to such disturbances may be particularly useful in linking mechanisms to disturbances and population responses (Williams et al., 2023). Linking individual- and population-level responses may be important to assessing whether OSW development drives significant adverse outcomes for shrimp fisheries.

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Table 3. Research questions for potential effects and pathways for effects of offshore wind (OSW) on taxonomic groups or fisheries in the Gulf of America (GoA). The taxonomic groups and fisheries include penaeid shrimp, reef fishes, coastal migratory pelagic (CMP) fishes, forage fishes, oceanic pelagic species, coastal elasmobranchs, and invasive species.

Taxa	Research questions
Penaeid shrimp	• Will shrimp habitat be affected, and at what scale, by direct disturbances from OSW, such as electric and magnetic fields (EMF), noise, and temperature? What are the physiological and behavioral mechanisms by which these direct disturbances may negatively affect shrimp? • How will benthic environmental conditions, including sediment composition and food availability, change through space and time and impact shrimp populations? • At what spatial scales of OSW development are shrimp fishing fleet behavior and profitability affected? What is the tipping point, regarding the number of wind farms and their spatial arrangement, for when the fleet needs to change how it operates? • Is it feasible for the shrimp fleet to supplement fishing-based income with collaborative efforts with researchers to collect in situ data relevant to assessing ecological OSW impacts?
Reef fishes	• To what extent does OSW infrastructure aggregate versus enhance productivity of reef fishes (i.e., growth, survival, reproduction)? How does the relative effect of aggregation versus production vary over space, time, and across reef fish taxa? • Will OSW serve as new spawning habitat for aggregating species? • How will OSW affect movement patterns, spatial distributions, and trophic ecology of reef fishes at local and seascape scales? • How will reef fish fisheries (commercial and recreational) target and exploit these new fish habitats? • What is the colonization trajectory of reef fishes at OSW sites? What factors affect the colonization trajectory (e.g., infrastructure spacing, depth, fishing pressure)? • How will sound, vibration, and EMF associated with OSW affect reef fishes? • How can OSW infrastructure, including scour protection and turbine foundations, be designed to maximize ecological benefits and minimize ecological risks to reef fishes? • How do reef fish community metrics (e.g., community composition, abundance, density, richness, diversity) on OSW sites compare with those at O&G, artificial reef, shipwreck, and natural reef sites (see Lemasson et al., 2024 for global meta-analysis)?
Coastal migratory pelagic (CMP) fishes	• Do the migratory patterns of CMP fishes currently coincide with planned OSW lease areas? Does the migratory behavior of CMP fishes change during different phases of OSW development (e.g., construction, operation, decommissioning)? • Will OSW change resource availability for CMP fishes via altered hydrodynamic regimes that affect pelagic food sources and floating structures (e.g., *Sargassum* spp.)? • To what extent does OSW infrastructure attract versus produce new biomass of CMP fishes? • What are the baseline levels of EMF, vibration, and sound within planned OSW lease areas? How do these conditions change with OSW development? How do any changes in these conditions affect CMP fishes related to their abundance, movement, temporal behavior, and other factors?
Forage fishes	• Could the presence of OSW in Louisiana state waters displace effort by the Gulf Menhaden *Brevoortia patronus* purse seine fishery? • Will OSW alter the spatial distribution and local densities of forage fishes? • Will OSW enhance production of forage fishes or aggregate them from nearby nonreef habitats?
Oceanic pelagic species	• To what degree do larvae of oceanic pelagic fishes interact with artificial structures and OSW?
Coastal elasmobranchs	• Will OSW increase interactions between humans and elasmobranchs, leading to novel or exacerbated effects? • Will EMF from OSW affect elasmobranch behavior and embryo development?
Invasive species	• Will OSW platforms serve as larval sources for Red Lionfish *Pterois volitans*, thereby exacerbating the Red Lionfish invasion? • Will OSW serve as stepping stones for invasive species range expansion? • What are the ecological impacts of Regal Damselfish *Neopomacentrus cyanomos* invasions, and will OSW mediate or exacerbate these impacts? • Will OSW serve as a critical habitat for fouling nonnative invertebrates? • Will OSW worsen jellyfish blooms, potentially disrupting fishing activity and/or forage fish that compete with jellyfish for planktonic food?

Reef fishes (Table 3)

In the GoA, reef fish communities are largely associated with hard-bottom, biogenic natural reefs or artificial structures. Small demersal reef fishes transfer energy via benthic and pelagic pathways to upper-trophic-level reef fishes, which support valuable recreational and commercial fisheries. Thirty-one GoA reef fish species undergo routine federal monitoring, assessment, and management (Gulf of Mexico Fishery Management Council [GMFMC], 2023). These are targeted by both recreational and commercial fisheries and are captured using a range of gears, including hook and line, vertical and bottom longlines, nets and traps, and spearfishing (data

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Table 4. Reef fishes in the Gulf of America that are important to recreational and commercial fisheries and their stock status as of December 31, 2023 (NMFS, 2024b; SEDAR, 2024). An empty status indicates that the species is not experiencing overfishing and is not overfished. Overfished indicates that the stock biomass is below the management target; experiencing overfishing indicates that fishing mortality is higher than the management target.

Reef fishes	Stock status (as of December 31, 2023)
Gag Grouper *Mycteroperca microlepis*	Overfished; experiencing overfishing
Gray Snapper *Lutjanus griseus*	
Gray Triggerfish *Balistes capriscus*	
Greater Amberjack *Seriola dumerili*	Overfished; experiencing overfishing
Lane Snapper *Lutjanus synagris*	Experiencing overfishing
Red Grouper *Epinephelus morio*	
Red Snapper *Lutjanus campechanus*	
Scamp *Mycteroperca phenax*	
Vermilion Snapper *Rhomboplites aurorubens*	
Yellowedge Grouper *Hyporthodus flavolimbatus*	Experiencing overfishing

summarized in Berenshtein et al., 2021). Some of the managed reef fish species have poor stock statuses (Table 4).

The effects of OSW on reef fishes will depend on the degree to which OSW structures function as artificial reefs, as reef fishes are strongly associated with hard-bottom habitat. Researchers are starting to examine how different types of OSW foundations and their components (e.g., scour protection), as well as their spatial configuration and density, affect fish abundance in other regions (Glarou et al., 2020; Werner et al., 2024), which could inform their functioning as artificial reefs. Although a global meta-analysis indicates that OSW structures may have higher fish abundances than adjacent soft-bottom habitats do, there is not enough existing evidence to detect the effects on fish abundance, biomass, or diversity relative to that of natural reefs (Lemasson et al., 2024). There are many additional knowledge gaps, including how trophic ecology, community metrics (e.g., community composition, diversity), and spatial distributions of reef fish (e.g., vertical and horizontal distribution, residence time, home range, etc.) near OSW infrastructure differ from those in existing GoA reef habitats. Finally, the potential ecological effects of OSW on reef fish will span population-to-ecosystem levels and must be considered cumulatively.

Reef fishes may also be affected by localized disturbances that are caused by OSW. For example, fish may respond to sound and vibration associated with preconstruction, construction, and operation of OSW (e.g., vessels, pile driving, turbine operation; reviewed in Popper et al., 2022). In addition, EMF that is associated with OSW infrastructure may affect reproduction, migration, and other behaviors in reef fish (Gill et al., 2020; Hutchison et al., 2020a). Developing rigorous monitoring programs, especially by implementing before–after–control–impact (BACI) or before–after-gradient (BAG) designs, will be critical to detecting and better understanding how reef fish respond specifically to OSW and related activities. It may also be possible to incorporate scour protection designs that are inspired by natural reefs and potentially other components associated with OSW turbine foundations to try to minimize the adverse effects (or optimize the beneficial effects) of OSW on reef fishes (Paxton et al., 2025).

Like other artificial structures, OSW infrastructure is expected to attract reef fish from nearby natural nonreef habitats (i.e., aggregation) and may enhance reef fish production as well (Bohnsack, 1989). Whether built structures ultimately yield net production gains is the topic of continued scientific research—that is, the "attraction–production" debate (e.g., Layman & Allgeier, 2020; Pickering & Whitmarsh, 1997). Evidence suggests that artificial reefs may enhance primary and/or secondary production (and therefore total reef fish biomass), although these effects are likely scale, place, and species specific (Claisse et al., 2014; Esquivel et al., 2022; Layman et al., 2016). Investigations in Europe, for example, demonstrate that small demersal, structure-associated fishes aggregate near OSW turbine foundations (Wilhelmsson et al., 2006), as do larger demersal piscivores (Bergstrom et al., 2013). However, enhanced production of reef fishes as a result of OSW has yet to be documented, but it will be critical to determine this for effective reef fish management (see Ecological processes for additional information).

Coastal migratory pelagic fishes (Table 3)

Coastal migratory pelagic (CMP) fishes in the GoA support important recreational and commercial fisheries, and there are some concerns over the health of these stocks. Gulf Cobia *Rachycentron canadum* are currently not overfished but are experiencing overfishing (NMFS, 2024b; SEDAR, 2020a). Although King Mackerel *Scomberomorus cavalla* and Spanish Mackerel *S. maculatus* were not estimated to be overfished or undergoing overfishing as of their last stock assessments (SEDAR, 2020b, 2023), individuals expressed concerns during recent GMFMC meetings about potential distributional shifts by these stocks in the GoA and changes in the availability of their prey. Dolphinfish *Coryphaena hippurus*, Wahoo *Acanthocybium solandri*, and Tripletail *Lobotes suirnamensis* also support fisheries, but they have not been assessed in the GoA. Stock assessments for CMP species in the GoA are made difficult by data limitations and by the fact that these species migrate across regional and national boundaries. As most CMP fishing is by hook and line near the surface, the development of OSW is unlikely to directly affect CMP fisheries, aside from the effects of artificial reefs or spatial exclusion management rules if they are implemented.

The reef effect of OSW development may have both positive and negative effects on CMP fish. We expect these effects to be less pronounced for CMP than those for reef fishes, given that CMP fishes spend time between fixed reef structures, floating structures (e.g., *Sargassum* spp., buoys, and floating inanimate objects), and the open ocean. Nevertheless, OSW structures are expected to provide habitat for larvae and juveniles (Hernandez et al., 2003; Lindquist et al., 2005), and turbine-induced wind wakes could increase productivity for earlier life stages of CMP species (Daewel et al., 2022). Adult CMP fishes typically rely on pelagic food sources (Finucane et al., 1990; Franks et al., 2008; Meyer & Franks, 1996; Oxenford & Hunte, 1999); however, OSW structures would provide high-relief habitat in the

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

entire water column that CMP fishes use for shelter, food supply, spawning, and spatial reference (reviewed in Franks, 2000; Sinopoli et al., 2017). The effects of OSW structure will likely be similar to the reef effects from O&G structures that aggregate CMP fishes: These include altered movement and migration patterns, aggregation of prey and changes in their food web linkages, and increased vulnerability to fishing (reviewed in Snodgrass et al., 2020). As with reef fishes, understanding the relative strength of aggregation (and increased vulnerability to exploitation) versus production gains remains an area of active research.

How physical disturbances, including EMF, vibration, and sound, from OSW may affect CMP fishes and their migratory behavior remains unknown, as does the degree to which CMP migratory patterns and the fisheries that target them overlap with lease areas. Some indirect effects may occur from de facto exclusions from active fishing gear in the areas. The combination of aggregation effects and the exclusion of shrimp trawling could reduce the bycatch of CMP fishes. However, fishing gear can become entangled in vertical and off-bottom structures and cause tear-offs or abrasions (Barnette, 2001); thus, fouled gear around OSW structures could potentially cause ghost fishing interactions. Spatial modeling analyses, using O&G infrastructure in the GoA as a proxy, could be used to explore the effects of aggregation, reduced risk of bycatch, and occurrence of fouled gear on CMP fishes.

Forage fishes (Table 3)

Forage species are typically small, short-lived (<5 years), schooling, or demersal fishes. They occupy an intermediate trophic level and are preyed upon by larger fish species of commercial and recreational importance, along with marine mammals and seabirds. Of the over 75 forage fish species that occur in the GoA (McEachran & Fechhelm, 1998, 2006), only a few of these species support fisheries. Gulf Menhaden are targeted by an industrial commercial purse-seine fishery that can remove the largest volume of catch in the USA GoA, which totaled 75% of all GoA commercial landings by mass in 2020 (NMFS, 2023). The Striped Mullet *Mugil cephalus* is another forage fish that is harvested commercially in the nearshore GoA, although only in Louisiana (Louisiana Department of Wildlife and Fisheries, 2020) and Florida (Florida Fish and Wildlife Conservation Commission, 2014). The only other forage fish that was historically commercially targeted and harvested in the region is Gulf Butterfish *Peprilus burti*, which is targeted using otter trawls that are pulled at depths greater than 100 m (Mareska & Bellais, 2023). However, recently, commercial interest in this species has declined (GMFMC, 2024, meeting minutes, 201, lines 29–38). The trophic roles of all forage fish and economic importance of the species that are specifically targeted by commercial fishing may drive interest in assessing interactions between forage fish and OSW.

The spatial overlap between planned OSW in the GoA and forage fishes is presently assumed to be minimal based on current distributions of fishing efforts, but it may increase in the future. The positioning of OSW structures in GoA federal waters is >16 km (9 nautical miles) from shore for Florida and Texas or 5 km (3 nautical miles) for Alabama, Mississippi, and Louisiana, and thus, it does not overlap with the habitat or fishing grounds of forage species. For example, the Gulf Menhaden fishery harvests most of its catch (93%) from within 16 km of the shoreline (Smith et al., 2002). However, spatial overlap may increase between OSW and forage fish habitats and fishing grounds with the addition of OSW leases within GoA state watersfor example, in Louisiana's 2023 lease sale (Baurick, 2023). Spatial overlap between GoA OSW and forage fishes may also increase if these species are driven to cooler, deeper waters in response to warming ocean temperatures (Pinsky et al., 2013, 2021). An initial concern of increased spatial overlap is the creation of de facto fishing exclusion zones, as purse seines and other harvest gear may be difficult to use around OSW infrastructure. Therefore, it is necessary to assess scenarios of how future OSW development in nearshore areas and climate change may alter interactions with forage species and fishery activity.

Because most forage fish species are not targeted by fisheries, the potential effects of OSW structures on forage fish have received less attention than potential effects on species that support commercial and recreational fisheries. Nevertheless, there are some studies concerning OSW interactions with forage fish in the northwest Atlantic. Friedland et al. (2023) documented the potential for significant spatial overlap between OSW and forage fishes, including Atlantic Menhaden, as multiple species preferred habitat within potential OSW lease areas. Additionally, Wilber et al. (2022a) found no effect of OSW on forage fish abundance (e.g., Atlantic Herring *Clupea harengus* and Atlantic Butterfish *Peprilus triacanthus* 7 years postconstruction. Although not specifically documented in response to OSW, forage fishes can be displaced by noise from simulated pile driving (Hawkins et al., 2014), ships (De Robertis & Handegard, 2013; Vabø et al., 2002), and seismic investigations (Slotte et al., 2004). This suggests that noise from OSW construction and operation may also influence the behavior of forage fishes, but more research is needed because there is limited evidence regarding behavioral and population-level responses of forage fishes to OSW.

The trophic importance of forage fishes, specifically Gulf Menhaden, in the GoA is becoming more established (Berenshtein et al., 2023; Sagarese et al., 2016, and references within). Therefore, future studies may consider assessing how OSW structures influence predator–prey interactions and the nutritional value of forage fish, as fish feeding behavior and physiology may change near OSW structures (e.g., Wilber et al., 2022b). It is also important to consider the responses of forage fish to OSW in the context of the attraction–production debate, as increased production of forage fish may lead to increased production or attraction of forage fish predators. Therefore, research is needed to assess how OSW may alter community-level interactions and trophic flows that are mediated by forage fish.

Oceanic pelagic fishes (Table 3)

Oceanic pelagic fishes in the GoA include teleost and elasmobranch species within the broad taxa of tunas (Scombridae), pelagic sharks (various families, e.g., Lamnidae, Carcharhinidae), billfishes (Istiophoridae), and Swordfish *Xiphias gladius*. Oceanic pelagic fishes also include highly migratory species that are characterized by extensive

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Table 5. Oceanic pelagic fishes in the Gulf of America that have stock statuses of concern and their stock status as of December 31, 2023 (NMFS, 2024b). Overfished indicates that the stock biomass is below the management target; experiencing overfishing indicates that fishing mortality is higher than the management target.

Oceanic pelagic fishes	Stock status (as of December 31, 2023)
Blue Marlin *Makaira nigricans*	Overfished; experiencing overfishing
White Marlin *Kajikia albida*	Overfished
Bigeye Tuna *Thunnus obesus*	Overfished
Dusky Shark *Carcharhinus obscurus*	Overfished; experiencing overfishing
Porbeagle Shark *Lamna nasus*	Overfished
Shortfin Mako Shark *Isurus oxyrinchus*	Overfished; experiencing overfishing

Table 6. Oceanic pelagic fishes in the Gulf of America that are prohibited from fisheries take and their status according to the International Union for Conservation of Nature (IUCN; Marshall et al., 2022; Pierce & Norman, 2016; Rigby et al., 2019).

Oceanic pelagic fishes	IUCN status
Giant Manta Ray *Mobula birostris*	Endangered
Oceanic Whitetip Shark *Carcharhinus longimanus*	Critically endangered
Whale Shark *Rhincodon typus*	Endangered

transoceanic movements and whose management necessitates international coordination. Many oceanic pelagic fishes have stock statuses of concern as of December 31, 2023 (NMFS, 2024b; Table 5). Some are prohibited from fisheries take and are on the International Union for Conservation of Nature Red List (Table 6). Given the current and expected locations of OSW development, we expect no or minimal interactions with Oceanic Whitetip Sharks *Carcharhinus longimanus*. Whale Sharks *Rhincodon typus* and Giant Manta Rays *Mobula birostris* could spatially overlap with OSW infrastructure, particularly in proposed development areas near the Flower Garden Banks National Marine Sanctuary (Stewart et al., 2018).

Siting options for OSW in the GoA are limited to shelf waters, and we expect them to have relatively minimal spatial overlap with most oceanic pelagic fishes that inhabit the open-ocean environment and typically occur offshore of continental shelf waters. This could change if future energy development extended farther offshore and employed the floating OSW infrastructure that is used in deepwater environments, in which case there are various concerns about its effects on these species, as summarized in Hendon et al. (2024). Offshore O&G structures affect pelagic longline fishing operations, which set main lines that can be 40 km long and drift for over 10 h. Such structures also attract both fish and fishermen, increasing vulnerability to fishing (Snodgrass et al., 2020). Although floating OSW is planned for other regions, the wide shelf area of the GoA provides more shallow-water leasing opportunities; also, floating OSW in the GoA is not expected within the near future. However, oceanic pelagic fishes may interact with OSW farms that are situated on the continental shelf because, similar to CMP, their larvae (e.g., Bluefin Tuna *Thunnus thynnus*) are found near O&G structures on the continental shelf (Hernandez et al., 2003). Continued telemetry tracking and spatial analyses will also be prudent for oceanic pelagic fishes given that climate change is expected to drive changing range distributions—for example, poleward shifts for Blue Marlin (Dale et al., 2022) and Bigeye Tuna (Erauskin-Extramiana et al., 2019).

Table 7. Coastal elasmobranchs, including inshore shark species and common northern Gulf of America batoids and, if applicable, their stock status as of December 31, 2023 (NMFS, 2024b). An empty status indicates the species is not experiencing overfishing and is not overfished. Overfished indicates that the stock biomass is below the management target; experiencing overfishing indicates that fishing mortality is higher than the management target.

Coastal elasmobranchs	Stock status (as of December 31, 2023)
Atlantic Sharpnose Shark *Rhizoprionodon terraenovae*	
Blacknose Shark *Carcharhinus acronotus*	Overfished; experiencing overfishing
Blacktip Shark *Carcharhinus limbatus*	
Bonnethead Shark *Sphyrna tiburo*	
Bull Shark *Carcharhinus leucas*	
Nurse Shark *Ginglymostoma cirratum*	
Sandbar Shark *Carcharhinus plumbeus*	Overfished
Sand Tiger Shark *Carcharias taurus*	
Scalloped Hammerhead Shark *Sphyrna lewini*	Overfished, experiencing overfishing
Southern Stingray *Dasyatis americana*	
Atlantic Stingray *Dasyatis sabina*	
Cownose Ray *Rhinoptera bonasus*	

Coastal elasmobranchs (Table 3)

Coastal elasmobranchs include inshore shark species and batoids, some of which have stock statuses of concern (Table 7). These species use coastal and nearshore habitats for various life stages, including nursery grounds, feeding, and some breeding activities (Chen, 2017, and references within). Their presence inshore makes them more accessible to coastal fisheries and increases their vulnerability to habitat degradation, bycatch mortality, and nearshore environmental stressors. Overexploitation has historically driven global declines in sharks and rays (Dulvy et al., 2014; MacNeil et al., 2020), which are particularly vulnerable due to their slow growth, late maturity, and production of few offspring. Such vulnerabilities and declines warrant strong consideration of the effects of OSW on coastal elasmobranchs.

Coastal elasmobranchs use electromagnetic-receptive sensory systems for many vital functions, such as foraging, predator avoidance, social communication, and navigation (Collin et al., 2015; Tricas & Sisneros, 2004). Therefore, specific research may be warranted to understand the interactions between elasmobranchs and EMFs produced by OSW. For

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

example, Hutchison et al. (2020c) found that EMFs caused a demonstrable increase in exploratory and foraging behaviors in Little Skate *Leucoraja erinacea*. These effects will likely be specific to the OSW installment (i.e., scale, output) and relate to the placement of subsea power cables, the strength of EMF they radiate, and whether the cables transmit alternating or direct current (Hermans et al., 2024). Some shark species undergo precise long-distance navigation using the Earth's magnetic field, produced by direct current (e.g., Keller et al., 2021). Tagging studies and spatial mapping analyses should examine behavioral changes in habitat usage and migratory patterns. A particular concern worth investigating is the potential disturbances by EMF on elasmobranch embryonic development (Hermans et al., 2024), as these could have population-level consequences. Such research should be species-specific, as effects will likely differ based on reproductive physiology (e.g., oviparous or viviparous species) and, for oviparous species, the specifics of where they lay their eggs.

Invasive species (Table 3)

If OSW infrastructure functions similarly to other artificial structures (e.g., artificial reefs, O&G structures), it could facilitate the expansion of invasive fishes and invertebrates in the GoA. The GoA currently has established invasions for two species of nonnative fishes of concern: Red Lionfish and Regal Damselfish. Red Lionfish represent the species of greatest concern given their direct demonstrated effects on prey species (e.g., Ballew et al., 2016; Dahl et al., 2017) and indirect effects on food webs (e.g., Chagaris et al., 2017, 2020; Tuttle, 2017). In the northern GoA, high-relief artificial structures serve as preferred Red Lionfish habitat and Red Lionfish densities on artificial structures can be 10–100 times those on natural reefs in the region (Dahl & Patterson, 2014 ; Harris et al., 2019). Offshore wind structures could serve as larval sources to enable the expansion from areas with higher densities of Red Lionfish—for example, from the Flower Gardens Banks National Marine Sanctuary (Blakeway et al., 2022) or eastern GoA reefs (Dahl & Patterson, 2020; Harris et al., 2023)—to reefs in the western GoA (Johnston et al., 2017). Nonnative Regal Damselfish were first observed on O&G and artificial structures in the northern GoA, and like Red Lionfish, they have higher densities on artificial structures than on natural reefs (Bennett, 2019). The highest densities are observed on relatively shallow artificial structures, which would have depths similar to those in the areas proposed for OSW development. Regal Damselfish were observed to comprise approximately half of the damselfish community in the north-central GoA (Tarnecki et al., 2021); however, their ecological impacts remain poorly understood and should be an area for further study. Collectively, the strong association between artificial structures and Red Lionfish, as well as uncertainties of Regal Damselfish invasions, warrant consideration for research concerning how these two species may interact with OSW structures.

Offshore wind and other artificial structures may serve as available habitat for nonnative fouling invertebrates. In the northern GoA, these have largely been observed on O&G platforms, with current invasions documented for nonnative orange cup corals *Tubastrea* spp., Australian spotted jellyfish *Phyllorhiza punctata* (during its polyp stage), the tunicate *Didemnum perlucidum*, and acorn barnacles (including triangle barnacle *Balanus trigonus*, striped barnacle *Amphibalanus amphitrite*, reticulated barnacle *A. reticulatus*, and titan acorn barnacle *Megabalanus coccopoma*) (Schulze et al., 2020). Although the overall community effects of these species are largely unknown, evidence indicates that cup corals disturb native communities of sessile invertebrates that may function as important habitat for fishery species (Lages et al., 2011). Additionally, blooms of the pelagic stage of Australian spotted jellyfish (among other jellyfish species) can cause large economic losses to shrimp fisheries by clogging trawl nets (Graham et al., 2003). Increased blooms of invasive jellyfish may also increase interspecific competition with planktivorous forage fish, as jellyfish and forage fish in the GoA may have strong diet overlap (D'Ambra et al., 2018). Future studies could assess the degree to which OSW may exacerbate interactions between invasive invertebrates and fisheries.

The role of OSW, O&G, and other artificial structures in facilitating biological invasions in the GoA remains an area of considerable uncertainty and active research (Schulze et al., 2020). Predicting the dynamics of invasive species remains notoriously difficult, as our understanding only comes after a species is established and the natural experiment of an invasion generally lacks control comparators (Strayer et al., 2017). Offshore wind platforms are expected to interact with invasive species in at least some capacity, and the relative interaction between invasive species and OSW versus O&G or other artificial structures will be important to evaluate, as there are already many artificial structures in the GoA (Figure 1D, 1E) and the initial expected footprint of OSW structures is comparatively small. Nevertheless, relevant pre- and post-OSW installation data collected at large temporal and spatial scales may elucidate these interactions, as there is a unique opportunity to "predict" and then validate the locations at which invasive species may arrive. Such data may allow for assessing the degree to which OSW platforms serve as stepping stones for range expansion by invasive species.

OVERARCHING OCEANOGRAPHIC AND ECOLOGICAL PROCESSES

Although different taxonomic groups require individualized attention, in terms of OSW effects, many (if not all) taxa will be affected by similar (if not the same) environmental processes. In this section, we consider broader oceanographic and ecological processes and how they may be affected by OSW. This is not intended to be a comprehensive review, as other studies have reviewed the effects of OSW on oceanographic and ecological processes (Gill et al., 2020; Reubens et al., 2014; van Berkel et al., 2020). Rather, we emphasize specific processes of interest to the GoA and provide context with representative examples from other regions (Table 8).

Oceanographic processes (Table 8)

Oceanographic processes and their inherent properties are important determinants of primary production, which supports secondary production of many fishery species via trophic pathways. Specifically, wind fields, stratification, and the effects of river discharge (hypoxia and turbidity), sediment

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Table 8. Research questions for potential effects and pathways for effects of offshore wind (OSW) on oceanographic and ecological processes in the Gulf of America.

Processes	Research questions
Oceanographic processes	• How will OSW affect wind fields and related physical oceanographic processes, such as turbulence, stratification, vertical/horizontal mixing, and upwelling/downwelling? • How will the physical effects of OSW affect biological processes and primary production? • Will OSW alter oxygen concentrations and at what temporal and spatial scales are those changes observable? • How will OSW affect benthic sediment dynamics, organic matter depositions, and thermal microhabitats, and will such changes affect benthic taxa, such as shrimp? • How do these oceanographic effects vary based on relative position to turbines (e.g., upwind versus downwind)?
Ecological processes	• To what degree will OSW enhance fish production, and will it be offset by aggregation from nearby habitats to OSW structures where they are more easily captured? • What are the successional dynamics of OSW structures, and how do they compare to O&G (or other artificial) structures and natural reefs? • How do fishes use the habitat created by OSW compared with O&G (or other artificial) structures and natural reefs? • Does the vertical relief of OSW turbines offer refuge from harmful environmental conditions, such as hypoxia? • How will OSW affect the physiology, behavior, and populations of reef fishes, forage fishes, and pelagic fishes? How will these interact with and influence trophic and community dynamics? • How will OSW affect connectivity and larval dispersal? • What are the fishery implications of changes in ecological processes?

transport, and deposition of particulate organic matter are often linked and important for fish and fisheries in the GoA (Bianchi et al., 2010; Coates et al., 2015; Hazen et al., 2009; Nagel et al., 2018). Because of their bottom-up effects on fish and fisheries, monitoring these abiotic, oceanographic processes will be important to help understand potential effects of OSW in the GoA.

Wind

The transport and retention of nutrient-rich surface waters that drive pelagic primary production is largely tied to wind, particularly in offshore waters where federal lease areas will occur (Muller-Karger et al., 2015). At local scales, changes in wind fields can lead to increased turbulence and mixing in turbine-adjacent waters (Carpenter et al., 2016). At larger scales, OSW farms can reduce wind stress within and downstream of OSW farms (referred to as a "wind wake") and can cause upwelling–downwelling dipoles (Broström, 2008; Christiansen & Hasager, 2005; van Berkel et al., 2020). The directionality and magnitude of primary production responses to such wind changes depend on location, depth, and scale. For example, at local scales, OSW may increase chlorophyll production and lead to chlorophyll aggregation, potentially due to increased upwelling and turbulence, but such signals may change at larger scales (Lu et al., 2022; van Berkel et al., 2020). To date, larger-scale (>100 km^2 from OSW farms) oceanographic effects from OSW have been evaluated for cool, nutrient-rich, North Atlantic waters (van Berkel et al., 2020), but further research is needed to understand its effects for subtropical systems, such as those in the GoA. Changes in wind and turbulence that are induced by OSW may also directly affect fish and fisheries through changes in the growth rates of larval fish due to turbulence-induced, altered predator–prey encounter rates (Kiørboe & MacKenzie, 1995). Fisheries ecologists and physical oceanographers should work together to coproduce study designs to monitor and detect these effects in the GoA.

Hypoxia and stratification

The GoA has the largest areas of hypoxia in the Western Hemisphere (Rabalais et al., 2001). The water column generally becomes highly stratified (unless mixed by tropical cyclones) during late summer, with lower water-column respiration depleting oxygen to levels that can form expansive hypoxic or anoxic areas that are hundreds of km^2 in size. Offshore wind structures can induce hypoxia due to increased respiration of higher levels of turbine-associated biomass, depending on spatial scale (Janßen et al., 2015). Stratification changes and thermocline shoaling induced by OSW can also directly affect the abundance and distribution of biota, including phytoplankton and zooplankton (Floeter et al., 2017). Mississippi River discharge largely affects hypoxia, stratification, and turbidity in the GoA. Offshore wind farms may increase or decrease turbidity, with such directionality changes potentially dependent on baseline conditions (van Berkel et al., 2020). In the GoA, it will be important for future work to separate the effects of OSW farms on the physical and chemical properties of the ecosystem from those of other drivers, such as the Mississippi River. These processes are very much intertwined, as OSW farms can simultaneously increase turbulence, nutrient transport, and chlorophyll production (Floeter et al., 2017).

Benthic and biogeochemical processes

Benthic oceanographic and biogeochemical processes may be disrupted by OSW construction (e.g., dredging) and operations (e.g., wind wakes; Coates et al., 2015; Nagel et al., 2018). For example, one study indicated that sediments directly adjacent to OSW scour protection zones exhibited lower grain sizes, potentially resulting in higher macrobenthos biomass (Coates et al., 2011). Biofouling of OSW turbines can also lead to organic enrichment of the benthos (Hutchison et al., 2020b). For example, biofouling mussels can produce strong fecal pellet rain that increases benthic organic matter deposition by 50% up to 5 km away from turbines (Ivanov et al., 2021). Thus, oceanographic effects are possible at relatively

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

large spatial scales, and biological hot spots may be generated by OSW invertebrate communities. Additionally, power cables from OSW farms may elevate the temperatures (~2.5°C) of the benthic boundary layer that is directly adjacent (within 1 m) to cables (Taormina et al., 2018). Although occurring at relatively small spatial scales, these effects may be important for smaller benthic fishery species, such as brown shrimp. Research assessing the effects of OSW on oceanographic and biogeochemical processes in the GoA will thus need to occur at variable scales, dimensions, and depths.

Ecological processes (Table 8)

Addition of new artificial hard-bottom and vertical habitat

The infrastructure for OSW has some similarities to existing artificial reefs and O&G platforms in the GoA, but it also has important differences that will influence our understanding of how OSW functions as an artificial hard-bottom and vertical habitat. Scour protection around OSW turbine foundations can be designed to function similarly to that of artificial reefs (Glarou et al., 2020). Also, OSW infrastructure is similar to O&G infrastructure in that it provides broad benthic cover and has a vertically extensive nature. Compared to monopile foundations, OSW jacket foundations, which have a lattice framework, will be especially similar to O&G platforms. Despite these similarities, OSW will differ from O&G in terms of spatial configuration and density. Although O&G platforms are irregularly arranged and spaced at varying distances apart, multiple OSW turbines will be placed at regular, gridded intervals within a lease area, separated by at least 1.85 km (1 nautical mile) for the purpose of navigational safety (U.S. Coast Guard, 2020). Because of the differences between OSW and existing human-made structures in the GoA, it is necessary to directly evaluate OSW as novel artificial habitat both at the seafloor and extending throughout the water column.

Given the foresight and the ability to collect data from pre- and postinstallation at appropriate spatial scales, OSW development presents a natural and large-scale opportunity to evaluate contributions from production and emigration to regional-scale fish populations. Carefully designing and implementing scientific studies (sensu Osenberg et al., 2002) could inform ecological questions, such as the attraction–production debate surrounding artificial structures and reef fishes. For example, greater relative abundance of Red Snapper has been observed on artificial structures (Gallaway et al., 2009), but whether this contributes to population-level changes remains debated and controversial (Cowan et al., 2011; Gardner et al., 2022; Karnauskas, Walter, et al., 2017; Szedlmayer & Shipp, 1994). By attracting fish, artificial structures can become hot spots for fishery removals by aggregating both fish and fishermen (i.e., ecological traps; Bohnsack, 1989; Gardner et al., 2022; Karnauskas, Walter, et al., 2017; Powers et al., 2018; Streich et al., 2017). Artificial structures may also increase production by providing additional settlement habitat (i.e., increased secondary production) for new recruits, thereby potentially increasing fishing opportunities (Bohnsack, 1989; Cowan et al., 2011). A large-scale, multidecade, and multiprong research and monitoring plan should use this opportunity to assess how OSW platforms affect fish production.

Succession

Biofouling organisms will rapidly colonize new hard substrate from OSW (Kerckhof et al., 2019). Over time, biofouling communities provide resources (e.g., food, habitat) for higher trophic levels. The composition of these biofouling communities may depend on construction materials (Petersen & Malm, 2006), timing of installation (Kerckhof et al., 2010, 2012), and location (Zupan et al., 2023). Benthic communities may also differ by the type of OSW turbines that are installed, as the type of foundation (jacket vs. monopile) and scour protection (rocks vs. sandbags) differ in surface area and structural complexity (Rumes et al., 2013). Biofouling communities on OSW turbines will go through successional stages, with early stages dominated by opportunistic species, intermediate stages being highly diverse, and climax stages dominated by a few species (e.g., Kerckhof et al., 2019). Climax communities on GoA O&G platforms and OSW infrastructure in the North Atlantic typically have high abundances of sponges, bryozoans, cnidarians (anemones, hydrozoans), and bivalves (Degraer et al., 2020; Gallaway & Lewbel, 1982). Full succession to a climax community can take at least 10 years for OSW communities (Kerckhof et al., 2019); thus, studies that are conducted at limited temporal spans may make misleading comparisons. Long-term monitoring will be required to assess the successional dynamics of benthic communities on OSW structures and to effectively compare OSW benthic communities with those found on other human-made structures (e.g., O&G, shipwrecks, artificial reefs) and natural reefs.

Habitat

Offshore wind structures often positively affect invertebrates and fishes (Galparsoro et al., 2022), but the quality of habitat that is provided by OSW infrastructure for a given species will depend on its use of the structure—for example, for foraging, shelter, navigation, or reproduction. Researchers who study feeding behavior found that fish obtain food resources from the hardened habitat created by OSW, but not all species use OSW as feeding grounds (reviewed in Hogan et al., 2023). Fish reproduction is less studied for OSW specifically, and findings are variable for O&G. Fish reproduction on O&G platforms was observed to be higher (Claisse et al., 2019), similar (Downey et al., 2018), or lower (Glenn et al., 2017) than that on natural reefs. Also, species may use the benthos-to-vertical structure of OSW to avoid environmental disturbances, such as moving higher up in the water column and avoiding hypoxic conditions (e.g., Carver, 2023; Reeves et al., 2018). Similarly, species vary in whether they remain present on or move from artificial structures during storm events, and it is hypothesized that fish are more likely to remain on larger, more stable structures, such as O&G and OSW structures during severe weather events, such as hurricanes (Topping & Szedlmayer, 2011). Overall, numerous metrics (e.g., diets, reproductive condition, movement, etc.) will need to be evaluated to obtain a comprehensive picture of the habitat quality of OSW for species and how it compares to that of natural reefs and other human-made structures in the GoA.

Habitat use of OSW structures will influence community dynamics (e.g., abundance, biomass, diversity). Demersal, structure-associated fishes and larger demersal piscivores can

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

aggregate near OSW turbine foundations (Bergstrom et al., 2013; Wilhelmsson et al., 2006). Generally, OSW increases adjacent fish abundance relative to reference sites (Lemasson et al., 2024; Methratta & Dardick, 2019), although this can be negated if fishing is allowed (Methratta & Dardick, 2019) and is inconclusive for fish biomass and diversity (Lemasson et al., 2024). Fish community structures around OSW will be influenced by intra- and interspecific interactions, which are typically difficult to study, especially for mobile species (Hogan et al., 2023). Therefore, it will be important to design research studies that can tease apart community interactions at OSW farms to provide a mechanistic understanding of fish community metrics (e.g., abundance, biomass, diversity).

Whether OSW habitat quality influences regional population dynamics of fishery species in the GoA will depend on the scale of OSW development. Even with over 20,000 known artificial reefs estimated to exist in the GoA (Gardner et al., 2022), artificial reefs cover a very small proportion (2%) of the total reef habitat, with the majority consisting of natural reefs (Gardner et al., 2022; Karnauskas, Walter et al., 2017). Also, researchers found that only 5% of GoA Red Snapper are located on artificial reefs (Stunz et al., 2021), only 1.2% of northern GoA Red Snapper biomass is found on O&G platforms (Osowski & Szedlmayer, 2022), and Red Snapper abundances on artificial structures only contribute 8% of the total population biomass (Karnauskas, Walter, et al., 2017). These studies suggest that OSW artificial habitat has a low probability of contributing to regionwide population-level increases in fish production. Nevertheless, OSW artificial habitat may function differently from other GoA artificial structures, so comparative (OSW versus O&G) and regionwide studies will be critical to elucidate the role of OSW in regional population dynamics of GoA fishery species.

Connectivity

Artificial structures can serve as connectivity corridors that fish use to move from one habitat to another within the seascape (Adams et al., 2014) and as stepping stones for larval dispersal of less motile taxa (Galaiduk et al., 2024). Whether this provides a net benefit to populations or the ecosystem remains uncertain (McLean et al., 2022). The potential positive effects of adding artificial reefs via OSW development include facilitating the movement of fish at their climate range edges (Paxton et al., 2019) or providing larval subsidies for nearshore reef fish as they undergo latitudinal migrations (Nishimoto et al., 2019). These connectivity corridors may, however, facilitate the spread of invasive species (see Invasive species [section 2] for further discussion of this topic).

Site fidelity to structured habitats may influence connectivity within populations, communities, and ecosystems (Lowerre-Barbieri et al., 2021; Patterson, 2007; Portnoy et al., 2022). Fish may exhibit high site fidelity to OSW farms (Reubens et al., 2013), O&G platforms (Everett et al., 2020), and other artificial reefs (Tharp et al., 2024; Topping & Szedlmayer, 2011). It is important to monitor a network of artificial structures when assessing fish movement, as single-point evaluations might falsely indicate low site fidelity to a single site when fish are in fact moving among various structured habitats within an area (e.g., Topping & Szedlmayer, 2011). Thus, the uniform spatial arrangement of OSW turbines within a lease area may be an interesting element that influences fish movement and site fidelity compared with arrangements in other artificial structures, which are typically more haphazardly distributed. It will also be important to monitor fish over long periods, as the presence and site fidelity of fish at OSW infrastructure can vary seasonally or due to spawning migrations (Reubens et al., 2013).

Offshore wind structures may also affect connectivity via larval dispersal. Fish movement between and site fidelity to OSW farms will influence larval dispersal, especially if OSW installations become spawning aggregation sites (Heyman et al., 2019; Perry & Heyman, 2020). Offshore wind structures can serve as sources of larvae for surrounding natural habitats, as modeled for O&G platforms in southern California (Nishimoto et al., 2019). However, larval movement patterns may be disrupted by changes in turbulence and stratification from OSW infrastructure (Ajmi et al., 2023; Chen et al., 2024). For example, simulation studies predict that OSW will displace larval dispersal and settlement of Atlantic sea scallops *Placopecten magellanicus* in the northeastern USA (Chen et al., 2024), in part due to enhanced turbulence and weakened stratification. Larval movement and survival are crucial for not only connectivity but also recruitment and other population-level effects (White et al., 2019, and references within). Larval movement, survival, and recruitment around OSW and surrounding natural and artificial habitats should be assessed to inform individual species' population dynamics and the functioning of the entire ecosystem. Understanding connectivity, site fidelity, and dispersal should constitute high-priority research topics, as they have direct implications for fisheries management.

IMPLICATIONS FOR MANAGEMENT AND HUMAN DIMENSIONS

Beyond anticipated physical, chemical, and biological effects of OSW maturation in the GoA, there will be potential effects on natural resource management and society. More than 15.8 million people live in U.S. coastal counties that are adjacent to the northern GoA (National Oceanic and Atmospheric Administration, 2023), and fishing is an important source of recreation and income. Here, we seek to identify the potential effects of OSW development on the human dimensions of the GoA, including fisheries management, fishing communities, and fisheries economies (Table 9).

Fisheries management (Table 9)

A concern accompanying OSW development is that it will affect scientific surveys that are used to monitor population trends. Using simulation analysis, Borsetti et al. (2023) found that excluding Atlantic surfclam *Spisula solidissima* assessment surveys from the regions designated for OSW development in the northeastern USA negatively affected the stock assessment reference points and increased uncertainty in estimates of spawning stock biomass and fishing mortality up to 17% and 7%, respectively. Standardized, long-term, fisheries-independent surveys track changes in abundance and population metrics over space and time for fish, invertebrates, and protected species (reviewed in Grüss et al., 2018) and provide critical data inputs for fisheries stock assessments. In the

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Table 9. Research questions for potential effects of offshore wind (OSW) development on human dimensions in the Gulf of America, including fisheries management, marine spatial planning (MSP), and social and economic effects.

Human dimensions	Research questions
Fisheries management	• How will the creation of OSW structures affect our ability to monitor and assess fisheries resources? • How will OSW change fishing behavior, in terms of number of trips, locations, and exploitation rates?
Marine spatial planning	• What fish and fisheries data layers are most critical for MSP, and how should these data layers be weighted relative to one another? • Can ecosystem models be used in MSP to account for dynamic ecological and fisheries interactions? • Can OSW be optimally co-sited with other ocean activities so that future ocean space is available for other ocean industries (e.g., aquaculture, tidal and wave energy, carbon capture, marine carbon dioxide removal, green hydrogen, transit, ecotourism)? • How should future OSW transmission and decommissioning be incorporated into the MSP process?
Social and economic effects	• What are the perceived and actual effects of OSW on the livelihoods of fishing communities, including commercial and recreational fisheries? • What are the potential effects on local economies if access to certain fishing grounds is lost or reduced? • What platforms and technologies can be leveraged to enable dialogue among scientists, managers, fishermen, and other resource users so that local concerns and knowledge properly inform research efforts and subsequent decision-making processes? • How do the social and economic impacts of OSW development vary across different sectors and communities at various timescales, and what options exist to mitigate them?

southeast Atlantic and GoA, a mitigation strategy was developed for fisheries-independent surveys (Hanisko et al., 2025). It will be important to minimize the effects of OSW development on fisheries-independent surveys and the metrics (e.g., mortality rates) that they produce (Haase et al., 2023).

The interplay between O&G infrastructure and the recreational fishing sector is a unique aspect of the GoA, as fishing activities are generally allowed on and immediately surrounding O&G platforms. Fishermen in the GoA have traditionally used O&G platforms as preferred fishing spots because they attract recreationally targeted species (Brashier, 1988; Stanley & Wilson, 1989). The use of O&G platforms in the GoA for fishing has been ubiquitous for over half a century and holds significant cultural importance (Dugas et al., 1979). Although existing and anticipated OSW development in the U.S. Northeast and Pacific is expected to restrict fishing activities around turbines, it is expected that OSW infrastructure generally will be open to fishing during operation in the GoA. The only expected restrictions are lack of access during the construction stage and that fishermen may be precluded from tying off to OSW structures. However, many anglers can lock on to sites using automated vessel positioning systems. As long as fishing access is not lost while OSW farms are operational, there has been general support for OSW development among charter captains that target reef fish (Klajbor et al., 2025). This general culture of openness to fishing on OSW infrastructure in the GoA presents novel considerations, research opportunities, and management implications. The potential effects of OSW development on economic and fishing behavior should also be considered. The addition of hardened benthic and vertical substrate may represent increased fishing opportunities for recreational and for-hire fishing operations. However, simulations for Atlantic surfclam fishermen in the northeastern USA predicted that OSW led to slightly increased fishing costs (<1–5%) and reduced revenues for fishing vessels and processors (~3–15%; Scheld et al., 2022). Also, OSW structures will present navigation hazards and will effectively exclude commercial trawlers from areas that have been fished historically, as discussed in previous sections. Broadly, different fisheries will be differentially affected such that the behavior and responses of various commercial and recreational fisheries should be continuously monitored as OSW development evolves in the GoA.

Marine spatial planning (Table 9)

Marine spatial planning (MSP) is a comprehensive, ecosystem-wide process that assesses new ocean uses and reconciles them with existing ocean uses and conservation of natural resources. In the GoA, regional MSP was conducted by the National Oceanic and Atmospheric Administration (NOAA) in partnership with BOEM to develop regional spatial models that identified areas of the ocean with the least spatial conflict and thus

Table 10. Submodel data categories and fisheries data layers incorporated into the marine spatial planning regional spatial models for evaluating the development of offshore wind (OSW) in the GoA (Randall et al., 2022).

Submodel data categories	Fisheries data layers
• Natural and cultural resources—e.g., sensitive habitats, protected resources • National security—e.g., military operation areas, special use airspace • Economics—e.g., OSW resource potential • Industry—e.g., vessel traffic, seafloor infrastructure, buoys • Logistics—e.g., distance from shore • Fisheries—see Fisheries data layers column • Constraints—e.g., military zones, coral, and hard bottom	• Commercial shrimp electronic logbook data • Highly migratory species pelagic longline gear observer data • Menhaden fishery data • Reef fish bandit gear fishing data • Reef fish longline gear fishing data • Southeast region headboat survey data

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

the highest opportunity for OSW development (Randall et al., 2022). The identified ocean areas (WEAs) were developed by aggregating 200 regional spatial data layers into a model. More specifically, each data layer was assigned to a submodel that was designated based on the data type (e.g., fisheries, industry, natural resources; Table 10). The seven resulting submodels were unweighted (i.e., the submodel from fisheries data had weight equal to that of the other six submodels of other ocean industries). Specific guidelines were also developed to resolve potential adverse effects on 23 species groups that are listed under the U.S. Endangered Species Act or the Marine Mammal Protection Act (Farmer et al., 2023). The final model applied a specific constraint for Rice's Whale *Balaenoptera ricei* and excluded the 100–400-m isobath.

Continued MSP research and communication will be critical to informing science-based wind energy development in the GoA, particularly within the wider context of changes in climate and ocean use. The current MSP tool does not incorporate ecosystem and fishery dynamics, which may lead to unanticipated consequences. Ecosystem modeling applications, such as Ecopath with Ecosim and Ecospace (EwEE), have been used for European OSW (Alexander et al., 2016; Püts et al., 2023) and MSP that may capture ecological feedback (Steenbeek et al., 2020). There are several EwEE models in the GoA that could be adapted to simulate MSP and OSW scenarios (Shaffer et al., 2023; Vilas et al., 2023). Ultimately, it will be necessary to balance wind energy development with both current (e.g., fisheries, O&G, mineral extraction, shipping, aquaculture) and potential future uses, such as tidal and wave renewable energy, marine carbon dioxide removal, carbon capture, and green hydrogen.

Social and economic effects (Table 9)

Although OSW in the GoA serves to address general societal values through energy independence and clean energy production, its development has the potential to disproportionately affect specific sectors and people. This may include immediate loss of fishing access and increased fuel costs to avoid traversing through OSW areas (Gray et al., 2016; Samoteskul et al., 2014), and disruptions to cultural ecosystem services like aesthetics, traditional heritage, and sense of place. As marine resource management alone fails to sufficiently address the equitability of such disruptions (Klain et al., 2014), the public should be engaged throughout the scientific process.

The primary effect of OSW on commercial fisheries will be the de facto exclusion of some active fishing gears (e.g., trawls) that will be unable to operate near OSW structures (Gray et al., 2005; Gray et al., 2016; Kirkpatrick et al., 2017; Mackinson et al., 2006), and the resulting loss of livelihood and fishing heritage (Alexander et al., 2013; Mackinson et al., 2006). Some pelagic gear fisheries may still be able to operate (Methratta et al., 2020). However, there are further concerns for potential crowding in remaining areas outside of OSW farms (e.g., Mackinson et al., 2006; ten Brink & Dalton, 2018), which may cause safety issues and increase interactions with protected species (Methratta et al., 2020; Schupp et al., 2021). Hook and line fishermen (both commercial and recreational) often support OSW development due to real or perceived abundance increases from the "artificial reef effect" of new structures (Methratta et al., 2020; Smythe et al., 2021; ten Brink & Dalton, 2018). In the northeastern USA, anglers have demonstrated general support for OSW development, but they have also stated that it detracts from the natural beauty and wilderness image of the sea (Bidwell et al., 2023; Hooper et al., 2017; Smythe et al., 2021).

Sense of place describes the meaning, or attachment, that a person or group ascribes to a specific geographical setting (Jorgensen & Stedman, 2001) and is a multidimensional concept consisting of feelings, associated memories, sensory perceptions, social connections, cultural rules, and the perceived relationship between self, others, and the place (Gustafson, 2001; Tuan, 1974; van Putten et al., 2018). Thus, sense of place is linked to a community's health and vulnerability to disturbances (Conley & Diamond, 2024). Place-based research where OSW development is already occurring globally reveals that communities may struggle to adjust to changes in their use of the ocean (Firestone et al., 2018), and individuals with strong place attachment tend to oppose OSW development (Bidwell, 2017). For example, the construction of the Block Island Wind Farm in the northeastern USA negatively affected residents' sense of place, even for individuals who supported the project (Russell et al., 2020), due to sentiment that the OSW turbines were unattractive, appeared industrial, and detracted from the overall character of the Rhode Island coast (Firestone et al., 2018; Russell et al., 2020). Sense of place is geographically specific; therefore, we expect that sentiments may differ in the GoA, as infrastructure developed for the O&G industry is already well established (Ditton & Auyong, 1984; Hooper et al., 2017). Studies of sense of place typically only commence after an adverse event has occurred. For example, coastal residents that depended on natural resources for their livelihoods (e.g., fisheries) experienced a loss of cultural heritage and a diminished sense of place and spiritual connection to the environment following the *Exxon Valdez* and Deepwater Horizon MC252 oil spills (Gill et al., 2014; Lee & Blanchard, 2012). Baseline data on sense of place and community health are needed (Sandifer et al., 2021). Such data, if available before OSW development, would enable developers and managers to better address the public's concerns and identify trade-offs (van Putten et al., 2018).

Engaging affected parties, including fishermen, OSW developers, and state and federal managers, early in the scientific process can align research priorities with societal values and address institutional biases (Voinov et al., 2014). Special efforts are needed to include underrepresented populations, such as racial, ethnic, or gender minorities and low-income groups who have often been disproportionately affected by and historically disenfranchised in environmental decision making (Bennett et al., 2023; Furman et al., 2023). To date, research on the social and economic effects of OSW on the commercial fishing industry and its communities remains relatively limited (Chaji & Werner, 2023). Participatory modeling can be employed in conjunction with other methodologies (e.g., ethnographic research and quantitative analysis of fishing effort and landings data) to elicit and integrate community knowledge to fill such gaps (Voinov et al., 2018)—for example, to support spatial planning in fisheries (Blake et al., 2017) and connect fishing effort data with shoreside communities (NOAA National

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Marine Fisheries Service, Southeast Fisheries Science Center, personal communication, 2024). The use of participatory modeling increases management transparency, reduces uncertainty (Röckmann et al., 2012), defines management objectives (Hobbs et al., 2002), and facilitates social learning (Li et al., 2016). Qualitative modeling tools can approximate scientific knowledge through crowdsourcing diverse perspectives, leveraging a "wisdom-of-crowds" approach for developing hypotheses or predictions (Aminpour et al., 2020). There are ample opportunities to apply such methods to leverage traditional and contemporary local ecological knowledge related to fisheries in the GoA to guide research directions that are related to OSW development.

DISCUSSION

The primary goals of OSW development in the USA are to produce decarbonized energy, advance national energy security, and support U.S. technologies and economies (The White House, 2021). In the GoA, a long history of industrial development has fostered a unique social, ecological, and regulatory environment among developers, managers, and members of the fishing industry, including commercial, recreational, and charter for-hire fishermen. Although BOEM has ongoing efforts to assess such effects, conducting additional and complementary research across all stages of OSW development is essential.

Disentangling OSW from concurrent environmental pressures

The GoA is already experiencing human and environmentally driven stressors and disturbances, such as ocean warming, ocean acidification, hypoxia, and harmful algal blooms. The introduction of OSW infrastructure has the potential to interact with these stressors and disturbances, and their interactions require further examination for responsible management and planning. There is little evidence suggesting potential alleviation of the adverse effects of climate change in the near future (Chagaris et al., 2019), and thus, it will be important to disentangle those effects from those driven by OSW development. Cumulative and interactive effects (e.g., additive, synergistic, antagonistic) of OSW development and other anthropogenic disruptions are expected and will need to be considered. For example, OSW structures support high biomasses of suspension-feeding invertebrate communities that increase the removal of organic matter from the water column, which is expected to increase more under ocean warming and acidification (Voet et al., 2022). These interactions may affect fishery species that rely on trophic pathways based on pelagic organic matter. Ultimately, the spatial and temporal extent of these interactions may determine the degree to which fisheries are affected, highlighting the need for regional scale, cumulative studies in the GoA.

Different perturbations are more prevalent in different regions of the GoA. For example, inner midshelf waters are more prone to hypoxia than offshore waters (Rabalais et al., 2001). Additionally, *Karenia brevis* or "red tide" blooms are more prevalent in southern GoA waters offshore Florida and Texas than in more northern and central GoA coastal states (e.g., Mississippi and Alabama; Soto et al., 2018). These spatial differences should be considered when evaluating the effects of forthcoming OSW structures and when disentangling these effects from those of other anthropogenic stressors. Relatedly, it will be important to disentangle and distinguish between the effects of OSW and other anthropogenic stressors to avoid mischaracterization of causal relationships. This issue is already present in the northeastern USA, where there is a public perception of linking OSW development to increased strandings of marine mammals, despite the current lack of a scientific link between OSW farms and these strandings (Thorne & Wiley, 2024).

Examining scale will be crucial for evaluating the effects of OSW. Although localized effects of OSW infrastructure on fish and fisheries can be studied at the turbine, cable, or even lease scale, regionally specific monitoring and science priorities are necessary due to the potential for cumulative or compounding effects (e.g., those associated with climate change) that cannot be identified at finer spatial and temporal scales (Methratta et al., 2023). An example of a program that addresses OSW at an ecosystem scale is NOAA's Gulf of Mexico Integrated Ecosystem Assessment (IEA) program, which summarizes and reports ecosystem-scale information in Ecosystem Status Reports (ESRs). These compile and generate ecosystem, biological, and social indicators, including climate drivers, physicochemical pressures, habitat changes, trophic states, ecosystem services, and human dimensions. To date, two ESRs for the GoA have been published by NOAA (Karnauskas, Keible, et al., 2017; Karnauskas et al., 2013). The data compiled for the ESRs enabled critical hypothesis testing on the linkages among drivers, pressures, and ecosystem states in the GoA (Karnauskas et al., 2015; Kilborn et al., 2018, 2024). Thus, the Integrated Ecosystem Assessment program and ESRs can inform empirically based assessments of concurrent environmental pressures in the GoA to help disentangle these from the effects of OSW farms.

Importance of baseline data

A central theme of the working group meeting was the importance of collecting comprehensive baseline data (biotic and abiotic variables) prior to the construction of OSW structures. Although there are several long-term monitoring surveys in the GoA (reviewed in Grüss et al., 2018), many of these surveys were designed to provide relative abundance indices for target marine resources across a wide spatial area for use in fisheries stock assessments. Therefore, most of these surveys do not have the statistical power to detect the effects of disturbances on finer-scale habitats (Powers et al., 2017). For example, the lack of baseline data prior to the Deepwater Horizon MC252 oil spill prevented a full assessment of the long-term effects on some marine resources and communities in the GoA (Powers et al., 2017), although population-level effects were more apparent for taxa with substantial prespill baseline community data (Patterson et al., 2023). The foresight from this working group can enable strategic plans for data collection in the proposed OSW lease areas before construction to help monitor responses using BACI or BAG sampling designs. Baseline data could be collected by supplementing ongoing monitoring surveys or by conducting new surveys that target specific habitats (e.g., benthic core grabs).

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Societal participation in the scientific process

In the GoA, science to inform federal fisheries management has a long history of meaningful public participation, not only in developing and vetting scientific products but also in public involvement in management advice based on this science. The U.S. Department of Commerce oversees this public participation through the Magnuson–Stevens Fishery Conservation and Management Act (1976). Specifically, the Magnuson–Stevens Fishery Conservation and Management Act created structures for public, interagency discussions, and decision-making processes for fisheries science and management in the USA with the establishment of eight Regional Fishery Management Councils, including the GMFMC. These councils include academic scientists, fishing industry members, and members of the public, and their long-standing existence and processes established expectations for transparency and engagement around fisheries issues (Wilson & McCay, 1998).

Alternatively, OSW development (specifically siting and permitting) is managed by the Department of the Interior under different legislation: the Outer Continental Shelf Lands Act (1953) and the Energy Policy Act (2005). These laws do not require a similar, comprehensive public engagement process for OSW decision making. Department of Interior agencies, such as BOEM and the Bureau of Safety and Environmental Enforcement, engage with fisheries constituents at their own discretion. To date, BOEM's GoA region has participated in council discussions, organized public outreach events, and held one-on-one meetings with fisheries constituents. However, because this is an ad hoc and voluntary process, as opposed to the Fishery Management Council system, the differences in how public input is gathered and used between the two resource management processes may exacerbate mistrust and hamper meaningful societal participation. Given the intersection between OSW, fisheries, and the GoA ecosystem, meaningful public participation throughout OSW development, installation, operations, and decommissioning is essential in the GoA and expected by the public. Improving public engagement in the GoA may be driven by individual states using their positions as decision makers to facilitate and support additional public participation processes. Additionally, regionwide public engagement could be supported by creating an independent regional science advisory body that includes agencies, scientists, the fishing community, and developers, similar to the Responsible Offshore Science Alliance and the Responsible Offshore Development Alliance in the Atlantic.

CONCLUSIONS

In the mid-20th century, O&G exploration began in federal and state waters of the GoA, altering the anthropogenic relationship with the region forever. The environmental impacts of this development became subject to assessment and consideration during the advent of national environmental policies like the National Environmental Policy Act of 1970. Although there are similarities between O&G and OSW development, it is wholly unlikely the O&G industry would have matured and developed in the GoA as it has if we had the scientific body of knowledge, regulatory environment, and assessment capabilities that we do now. This speculation, considered in the context of OSW development in the same region, begs the question: What would we have done differently if we knew then what we know now about the potential effects of offshore energy development? This type of opportunity in which society can anticipate and plan for a major intrusion into the natural environment with years of advance notice is unprecedented and underscores the importance of codeveloping scientific and monitoring priorities.

The timeline of OSW development provides scientific and natural resource management communities in the GoA region with a unique opportunity to shape the course of OSW development. Such efforts should encourage novel and applied science and sustainable development opportunities while mitigating for known (and potential) adverse effects and fostering positive effects, thereby maximizing sustainable ecological and economic benefits for all sectors and communities involved. Such research could help inform responsible and equitable construction, operation, and decommissioning of OSW infrastructure (precradle to postgrave) to support the coexistence of environmental and economic prosperity.

DATA AVAILABILITY

There are no new data associated with this article.

ETHICS STATEMENT

This research met the ethical guidelines and legal requirements of the country in which it was performed.

FUNDING

This article is a result of research that was funded by the National Oceanic and Atmospheric Administration's RESTORE Science Program (ROR—https://ror.org/0042xzm63) under award NA22NOS4510203 to the National Center for Ecological Analysis and Synthesis at the University of California, Santa Barbara as part of the Impacts of Offshore Wind on Gulf of Mexico Fish and Fisheries of the Gulf Ecosystem Initiative.

CONFLICTS OF INTEREST

None declared.

ACKNOWLEDGMENTS

We thank Mariana Steen (BOEM) for her participation in our working group and for the valuable insights, knowledge, and feedback she provided for this article.

REFERENCES

Adams, T. P., Miller, R. G., Aleynik, D., & Burrows, M. T. (2014). Offshore marine renewable energy devices as stepping stones across biogeographic boundaries. *Journal of Applied Ecology*, *51*, 330–338. https://doi.org/10.1111/1365-2664.12207

Ajmi, S., Boutet, M., Bennis, A. C., Dauvin, J. C., & Pezy, J. P. (2023). Numerical study of turbulent wake of offshore wind turbines and retention time of larval dispersion. *Journal of Marine Science and Engineering*, *11*, Article 2152. https://doi.org/10.3390/jmse11112152

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Alexander, K. A., Meyjes, S. A., & Heymans, J. J. (2016). Spatial ecosystem modelling of marine renewable energy installations: Gauging the utility of ecospace. *Ecological Modelling*, *331*, 115–128. https://doi.org/10.1016/j.ecolmodel.2016.01.016

Alexander, K. A., Potts, T., & Wilding, T. A. (2013). Marine renewable energy and Scottish west coast fishers: Exploring impacts, opportunities and potential mitigation. *Ocean & Coastal Management*, *75*, 1–10. https://doi.org/10.1016/j.ocecoaman.2013.01.005

Aminpour, P., Gray, S. A., Jetter, A. J., Introne, J. E., Singer, A., & Arlinghaus, R. (2020). Wisdom of stakeholder crowds in complex social–ecological systems. *Nature Sustainability*, *3*, 191–199. https://doi.org/10.1038/s41893-019-0467-z

Arreguín-Sánchez, F., del Monte-Luna, P., & Zetina-Rejón, M. J. (2015). Climate change effects on aquatic ecosystems and the challenge for fishery management: Pink shrimp of the southern Gulf of Mexico. *Fisheries*, *40*, 15–19. https://doi.org/10.1080/03632415.2015.988075

Ballew, N. G., Bacheler, N. M., Kellison, G. T., & Schueller, A. M. (2016). Invasive lionfish reduce native fish abundance on a regional scale. *Scientific Reports*, *6*, Article 32169. https://doi.org/10.1038/srep32169

Barnette, M. C. (2001). *A review of the fishing gear utilized within the Southeast Region and their potential impacts on essential fish habitat* (Technical Memorandum NMFS-SEF SC-44 9). National Oceanic and Atmospheric Administration. https://repository.library.noaa.gov/view/noaa/8527

Baurick, T. (2023, December 13). Louisiana signs agreements to build first offshore wind farms in state waters. NOLA.com. https://www.nola.com/news/environment/louisiana-signs-agreements-for-its-first-offshore-wind-farms/article_1f2a8708-99f2-11ee-a5c8-976e6eb24217.html

Beier, P., Hansen, L. J., Helbrecht, L., & Behar, D. (2017). A how-to guide for coproduction of actionable science. *Conservation Letters*, *10*, 288–296. https://doi.org/10.1111/conl.12300

Bennett, C. (2019). First record of the non-indigenous Indo-Pacific damselfish, *Neopomacentrus cyanomos* (Bleeker, 1856) in the northern Gulf of Mexico. *BioInvasions Records*, *8*, 154–166. https://doi.org/10.3391/bir.2019.8.1.17

Bennett, N. J., Alava, J. J., Ferguson, C. E., Blythe, J., Morgera, E., Boyd, D., & Côté, I. M. (2023). Environmental (in)justice in the Anthropocene ocean. *Marine Policy*, *147*, Article 105383. https://doi.org/10.1016/j.marpol.2022.105383

Berenshtein, I., Sagarese, S. R., Lauretta, M. V., & Chagaris, D. D. (2021). *Technical documentation of a U.S. Gulf of Mexico-wide Ecosystem model* (Technical Memorandum NMFS-SEFSC-751). National Oceanic and Atmospheric Administration. https://doi.org/10.25923/zj8t-e656

Berenshtein, I., Sagarese, S. R., Lauretta, M. V., Schueller, A. M., & Chagaris, D. D. (2023). Identifying trade-offs and reference points in support of ecosystem approaches to managing Gulf of Mexico menhaden. *Frontiers in Marine Science*, *9*, Article 935324. https://doi.org/10.3389/fmars.2022.935324

Bergström, L., Sundqvist, F., & Bergström, U. (2013). Effects of an offshore wind farm on temporal and spatial patterns in the demersal fish community. *Marine Ecology Progress Series*, *485*, 199–210. https://doi.org/10.3354/meps10344

Bianchi, T. S., DiMarco, S. F., Cowan, J. H., Jr., Hetland, R. D., Chapman, P., Day, J. W., & Allison, M. A. (2010). The science of hypoxia in the northern Gulf of Mexico: A review. *The Science of the Total Environment*, *408*, 1471–1484. https://doi.org/10.1016/j.scitotenv.2009.11.047

Bidwell, D. (2017). Ocean beliefs and support for an offshore wind energy project. *Ocean & Coastal Management*, *146*, 99–108. https://doi.org/10.1016/j.ocecoaman.2017.06.012

Bidwell, D., Smythe, T., & Tyler, G. (2023). Anglers' support for an offshore wind farm: Fishing effects or clean energy symbolism. *Marine Policy*, *151*, Article 105568. https://doi.org/10.1016/j.marpol.2023.105568

Blake, D., Augé, A. A., & Sherren, K. (2017). Participatory mapping to elicit cultural coastal values for marine spatial planning in a remote archipelago. *Ocean & Coastal Management*, *148*, 195–203. https://doi.org/10.1016/j.ocecoaman.2017.08.010

Blakeway, R. D., Fogg, A. Q., Johnston, M. A., Rooker, J. R., & Jones, G. A. (2022). Key life history attributes and removal efforts of invasive lionfish (*Pterois volitans*) in the Flower Garden Banks National Marine Sanctuary, northwestern Gulf of Mexico. *Frontiers in Marine Science*, *9*, Article 774407. https://doi.org/10.3389/fmars.2022.774407

Bohnsack, J. A. (1989). Are high densities of fishes at artificial reefs the result of habitat limitation or behavioral preference? *Bulletin of Marine Science*, *44*, 631–645.

Borsetti, S., Munroe, D. M., Scheld, A. M., Powell, E. N., Klinck, J. M., & Hofmann, E. E. (2023). Potential repercussions of offshore wind energy development in the northeast United States for the Atlantic surfclam survey and population assessment. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science*3, *15*, Article e10228. https://doi.org/10.1002/mcf2.10228

Brashier, J. (1988). Coexistence of fishing and oil and gas industries in the Gulf of Mexico. In *Proceedings of OCEANS'88. "A Partnership of Marine Interests"* (pp. 136–141). IEEE. https://doi.org/10.1109/OCEANS.1988.794841

Broström, G. (2008). On the influence of large wind farms on the upper ocean circulation. *Journal of Marine Systems*, *74*, 585–591. https://doi.org/10.1016/j.jmarsys.2008.05.001

Brunik, K., Thomas, J. J., Clark, C. E., Duffy, P., Kotarbinski, M., Kee, J., Grant, E., Starke, G. M., Riccobono, N., Koleva, M. N., Reznicek, E. P., Barker, A., & King, J. (2024). Potential for large-scale deployment of offshore wind-to-hydrogen systems in the United States. *Journal of Physics: Conference Series*, *2767*, Article 062017. https://doi.org/10.1088/1742-6596/2767/6/062017

Bugnot, A. B., Mayer-Pinto, M., Airoldi, L., Heery, E. C., Johnston, E. L., Critchley, L. P., Strain, E. M. A., Morris, R. L., Loke, L. H. L., Bishop, M. J., Sheehan, E. V., Coleman, R. A., & Dafforn, K. A. (2021). Current and projected global extent of marine built structures. *Nature Sustainability*, *4*, 33–41. https://doi.org/10.1038/s41893-020-00595-1

Bureau of Ocean Energy Management. (2024). *The renewable energy process: Leasing to operations* [Fact sheet]. Retrieved May, 2024, from https://www.boem.gov/sites/default/files/documents/renewable-energy/state-activities/RE-Leasing-Process-Poster.pdf

Carpenter, J. R., Merckelbach, L., Callies, U., Clark, S., Gaslikova, L., & Baschek, B. (2016). Potential impacts of offshore wind farms on north sea stratification. *PLoS One*, *11*, Article e0160830. https://doi.org/10.1371/journal.pone.0160830

Carver, J. (2023). *Reef fish movement dynamics and recruitment to artificial habitat* [Master's thesis, Louisiana State University]. Louisiana State University Repository. https://repository.lsu.edu/gradschool_theses/5814

Chagaris, D., Binion-Rock, S., Bogdanoff, A., Dahl, K., Granneman, J., Harris, H., Mohan, J., Rudd, M. B., Swenarton, M. K., Ahrens, R., & Patterson, W. F., III. (2017). An ecosystem-based approach to evaluating impacts and management of invasive lionfish. *Fisheries*, *42*, 421–431. https://doi.org/10.1080/03632415.2017.1340273

Chagaris, D. D., Patterson, W. F., & Allen, M. S. (2020). Relative effects of multiple stressors on reef food webs in the northern Gulf of Mexico revealed via ecosystem modeling. *Frontiers in Marine Science*, *7*, Article 00513. https://doi.org/10.3389/fmars.2020.00513

Chagaris, D., Sagarese, S., Farmer, N., Mahmoudi, B., de Mutsert, K., VanderKooy, S., Patterson, W. F., III, Kilgour, M., Schueller, A., Ahrens, R., & Lauretta, M. (2019). Management challenges are opportunities for fisheries ecosystem models in the Gulf of Mexico. *Marine Policy*, *101*, 1–7. https://doi.org/10.1016/j.marpol.2018.11.033

Chaji, M., & Werner, S. (2023). Economic impacts of offshore wind farms on fishing industries: Perspectives, methods, and knowledge gaps. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science*, *15*, Article e10237. https://doi.org/10.1002/mcf2.10237

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Chen, Y. (2017). Fish resources of the Gulf of Mexico. In C. Herb Ward (Ed.), *Habitats and biota of the Gulf of Mexico: Before the Deepwater Horizon Oil Spill. Volume 2: Fish resources, fisheries, sea turtles, avian resources, marine mammals, diseases and mortalities* (pp. 869–1038). Springer. https://link.springer.com/chapter/10.1007/978-1-4939-3456-0_1

Chen, C., Zhao, L., Lin, H., He, P., Li, S., Wu, Z., Qi, J., Xu, Q., Stokesbury, K., & Wang, L. (2024). Potential impacts of offshore wind energy development on physical processes and scallop larval dispersal over the US northeast shelf. *Progress in Oceanography*, *224*, Article 103263. https://doi.org/10.1016/j.pocean.2024.103263

Christiansen, M. B., & Hasager, C. B. (2005). Wake effects of large offshore wind farms identified from satellite SAR. *Remote Sensing of Environment*, *98*, 251–268. https://doi.org/10.1016/j.rse.2005.07.009

Claisse, J. T., Love, M. S., Meyer-Gutbrod, E. L., Williams, C. M., & Pondella, D. J., II. (2019). Fishes with high reproductive output potential on California offshore oil and gas platforms. *Bulletin of Marine Science*, *95*, 515–534. https://doi.org/10.5343/bms.2019.0016

Claisse, J. T., Pondella, D. J., Love, M., Zahn, L. A., Williams, C. M., Williams, J. P., & Bull, A. S. (2014). Oil platforms off California are among the most productive marine fish habitats globally. *Proceedings of the National Academy of Sciences of the United States of America*, *111*, 15462–15467. https://doi.org/10.1073/pnas.1411477111

Coates, D., Vanaverbeke, J., Rabaut, M., & Vincx, M. (2011). Soft-sediment macrobenthos around offshore wind turbines in the Belgian part of the north sea reveals a clear shift in species composition. In S. Degraer, R. Brabant, & B. Rumes (Eds.), *Offshore wind farms in the Belgian part of the north sea: Selected findings from the baseline and targeted monitoring* (pp. 47–63). Royal Belgian Institute of Natural Sciences.

Coates, D. A., Van Hoey, G., Colson, L., Vincx, M., & Vanaverbeke, J. (2015). Rapid macrobenthic recovery after dredging activities in an offshore wind farm in the Belgian part of the north sea. *Hydrobiologia*, *756*, 3–18. https://doi.org/10.1007/s10750-014-2103-2

Collin, S. P., Kempster, R. M., & Yopak, K. E. (2015). How elasmobranchs sense their environment. In R. E. Shadwick, A. P. Farrell, & C. J. Brauner (Eds.), *Physiology of elasmobranch fishes: Structure and interaction with environment* (Fish Physiology 34A, pp. 19–99). Academic Press.

Conley, E. D., & Diamond, E. P. (2024). Angler's sense of place as an indicator for perceived vulnerability to shifting stock distributions. *Marine Policy*, *161*, Article 106007. https://doi.org/10.1016/j.marpol.2024.106007

Cook, H. L., & Lindner, M. J. (1970). *Synopsis of biological data on the brown shrimp,* Penaeus aztecus *Ives, 1891* (Fisheries Synopsis 102). Food and Agriculture Organization of the United Nations.

Cowan, J. H., Grimes, C. B., Patterson, W. F., III, Walters, C. J., Jones, A. C., Lindberg, W. J., Sheehy, D. J., Pine, W. E., Powers, J. E., Campbell, M. D., & Lindeman, K. C. (2011). Red Snapper management in the Gulf of Mexico: Science-or faith-based? *Reviews in Fish Biology and Fisheries*, *21*, 187–204. https://doi.org/10.1007/s11160-010-9165-7

Daewel, U., Akhtar, N., Christiansen, N., & Schrum, C. (2022). Offshore wind farms are projected to impact primary production and bottom water deoxygenation in the north sea. *Communications Earth & Environment*, *3*, Article 292. https://doi.org/10.1038/s43247-022-00625-0

Dahl, K. A., & Patterson, W. F., III. (2014). Habitat-specific density and diet of rapidly expanding invasive Red Lionfish, *Pterois volitans*, populations in the northern Gulf of Mexico. *PLoS One*, *9*, Article e105852. https://doi.org/10.1371/journal.pone.0105852

Dahl, K. A., & Patterson, W. F., III. (2020). Movement, home range, and depredation of invasive Lionfish revealed by fine-scale acoustic telemetry in the northern Gulf of Mexico. *Marine Biology*, *167*, Article 111. https://doi.org/10.1007/s00227-020-03728-4

Dahl, K. A., Patterson, W. F., III, Robertson, A., & Ortmann, A. C. (2017). DNA barcoding significantly improves resolution of invasive Lionfish diet in the northern Gulf of Mexico. *Biological Invasions*, *19*, 1917–1933. https://doi.org/10.1007/s10530-017-1407-3

Dale, J. J., Brodie, S., Carlisle, A. B., Castleton, M., Hazen, E. L., Bograd, S. J., & Block, B. A. (2022). Global habitat loss of a highly migratory predator, the Blue Marlin (*Makaira nigricans*). *Diversity & Distributions*, *28*, 2020–2034. https://doi.org/10.1111/ddi.13606

D'Ambra, I. D., Graham, W. M., Carmichael, R. H., & Hernandez, F. J., Jr. (2018). Dietary overlap between jellyfish and forage fish in the northern Gulf of Mexico. *Marine Ecology Progress Series*, *587*, 31–40. https://doi.org/10.3354/meps12419

Degraer, S., Carey, D. A., Coolen, J. W., Hutchison, Z. L., Kerckhof, F., Rumes, B., & Vanaverbeke, J. (2020). Offshore wind farm artificial reefs affect ecosystem structure and functioning. *Oceanography*, *33*, 48–57. https://doi.org/10.5670/oceanog.2020.405

De Robertis, A., & Handegard, N. O. (2013). Fish avoidance of research vessels and the efficacy of noise-reduced vessels: A review. *ICES Journal of Marine Science*, *70*, 34–45. https://doi.org/10.1093/icesjms/fss155

Ditton, R. B., & Auyong, J. (1984). *Fishing offshore platforms, central Gulf of Mexico: An analysis of recreational and commercial fishing use at 164 major offshore petroleum structures*. U.S. Department of the Interior, Minerals Management Service.

Doty, H. (2023, August 24). *What does offshore wind energy look like today?* [Blog]. Office of Energy Efficiency and Renewable Energy, Wind Energy Technologies Office. https://www.energy.gov/eere/wind/articles/what-does-offshore-wind-energy-look-today

Downey, C. H., Streich, M. K., Brewton, R. A., Ajemian, M. J., Wetz, J. J., & Stunz, G. W. (2018). Habitat-specific reproductive potential of Red Snapper: A comparison of artificial and natural reefs in the western Gulf of Mexico. *Transactions of the American Fisheries Society*, *147*, 1030–1041. https://doi.org/10.1002/tafs.10104

Dugas, R., Guillory, V., & Fischer, M. (1979). Oil rigs and offshore sport fishing in Louisiana. *Fisheries*, *4*, 2–10. https://doi.org/10.1577/1548-8446-4-6

Dulvy, N. K., Fowler, S. L., Musick, J. A., Cavanagh, R. D., Kyne, P. M., Harrison, L. R., Carlson, J. K., Davidson, L. N., Fordham, S. V., Francis, M. P., Pollock, C. M., Simpfendorfer, C. A., Burgess, G. H., Carpenter, K. E., Compagno, L. J., Ebert, D. A., Gibson, C., Heupel, M. R., Livingstone, S. R., … White, W. T. (2014). Extinction risk and conservation of the world's sharks and rays. *eLife*, *3*, Article e00590. https://doi.org/10.7554/eLife.00590

Energy Policy Act. (2005). 42 U.S.C. § 13201 *et seq*.

Erauskin-Extramiana, M., Arrizabalaga, H., Hobday, A. J., Cabré, A., Ibaibarriaga, L., Arregui, I., Murua, H., & Chust, G. (2019). Large-scale distribution of tuna species in a warming ocean. *Global Change Biology*, *25*, 2043–2060. https://doi.org/10.1111/gcb.14630

Esquivel, K. E., Hesselbarth, M. H. K., & Allgeier, J. E. (2022). Mechanistic support for increased primary production around artificial reefs. *Ecological Applications*, *32*, Article e2617. https://doi.org/10.1002/eap.2617

Everett, A., Szedlmayer, S., & Gallaway, B. (2020). Movement patterns of Red Snapper *Lutjanus campechanus* based on acoustic telemetry around oil and gas platforms in the northern Gulf of Mexico. *Marine Ecology Progress Series*, *649*, 155–173. https://doi.org/10.3354/meps13448

Farmer, N. A., Garrison, L. P., Litz, J. A., Ortega-Ortiz, J. G., Rappucci, G., Richards, P. M., Powell, J. R., Bethea, D. M., Jossart, J. A., & Randall, A. L. (2023). Protected species considerations for ocean planning: A case study for offshore wind energy development in the US Gulf of Mexico. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science*, *15*, Article e10246. https://doi.org/10.1002/mcf2.10246

Finucane, J. H., Grimes, C. B., & Naughton, S. P. (1990). Diets of young King and Spanish mackerel off the southeast United States. *Northeast Gulf Science*, *11*, 145–153. https://doi.org/10.18785/negs.1102.07

Firestone, J., Bidwell, D., Gardner, M., & Knapp, L. (2018). Wind in the sails or choppy seas? People-place relations, aesthetics and public support for the United States' first offshore wind project. *Energy Research & Social Science*, *40*, 232–243. https://doi.org/10.1016/j.erss.2018.02.017

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Floeter, J., van Beusekom, J. E., Auch, D., Callies, U., Carpenter, J., Dudeck, T., Eberle, S., Eckhardt, A., Gloe, D., Hänselmann, K., & Hufnagl, M. (2017). Pelagic effects of offshore wind farm foundations in the stratified north sea. *Progress in Oceanography*, *156*, 154–173. https://doi.org/10.1016/j.pocean.2017.07.003

Florida Fish and Wildlife Conservation Commission. (2014). *The 2014 stock assessment update for Striped Mullet,* Mugil cephalus, *in Florida* [FWC IHR-2014-065]. https://myfwc.com/media/13333/stripedmullet-assessment-2014-final.pdf

Franks, J. (2000). A review: Pelagic fishes at petroleum platforms in the northern Gulf of Mexico; diversity, interrelationships, and perspective. In J.-Y. Le Gall, P. Cayre, & M. Taquet (Eds.), *Pêche thonière et dispositifs de concentration de poissons* (pp. 502–515). Ifremer. https://archimer.ifremer.fr/doc/00042/15301/

Franks, J. S., Hoffmayer, E. R., Ballard, J. R., Garber, N. M., & Garber, A. F. (2008). Diet of Wahoo, *Acanthocybium solandri*, from the northcentral Gulf of Mexico. *Proceedings of the Gulf and Caribbean Fisheries Institute*, *60*, 353–362.

Friedland, K. D., Adams, E. M., Goetsch, C., Gulka, J., Brady, D. C., Rzeszowski, E., Crear, D. P., Gaichas, S., Gill, A. B., McManus, M. C., & Methratta, E. T. (2023). Forage fish species prefer habitat within designated offshore wind energy areas in the U.S. northeast shelf ecosystem. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science*, *15*, Article e10230. https://doi.org/10.1002/mcf2.10230

Furman, K. L., Harlan, S. L., Barbieri, L., & Scyphers, S. B. (2023). Social equity in shore-based fisheries: Identifying and understanding barriers to access. *Marine Policy*, *148*, Article 105355. https://doi.org/10.1016/j.marpol.2022.105355

Galaiduk, R., McLean, D. L., Speed, C. W., Greer, D., McIntosh, R., & Treml, E. A. (2024). Offshore oil and gas infrastructure plays a minor role in marine metapopulation dynamics. *The Science of the Total Environment*, *934*, Article 172981. https://doi.org/10.1016/j.scitotenv.2024.172981

Gallaway, B. J., & Lewbel, G. S. (1982). *The ecology of petroleum platforms in the northwestern Gulf of Mexico: A community profile* (FWS/OBS 82/27). U.S. Fish and Wildlife Service.

Gallaway, B. J., Raborn, S. W., McCain, K. A., Beyea, R. T., Dufault, S., Heyman, W., Putman, N. F., & Egerton, J. (2021). Absolute abundance estimates for Red Snapper, Greater Amberjack, and other federally managed fish on offshore petroleum platforms in the Gulf of Mexico. *North American Journal of Fisheries Management*, *41*, 1665–1690. https://doi.org/10.1002/nafm.10678

Gallaway, B. J., Szedlmayer, S. T., & Gazey, W. J. (2009). A life history review for Red Snapper in the Gulf of Mexico with an evaluation of the importance of offshore petroleum platforms and other artificial reefs. *Reviews in Fisheries Science*, *17*, 48–67. https://doi.org/10.1080/10641260802160717

Galparsoro, I., Menchaca, I., Garmendia, J. M., Borja, Á, Maldonado, A. D., Iglesias, G., & Bald, J. (2022). Reviewing the ecological impacts of offshore wind farms. *npj Ocean Sustainability*, *1*, Article 1. https://doi.org/10.1038/s44183-022-00003-5

Gardner, C., Goethel, D. R., Karnauskas, M., Smith, M. W., Perruso, L., & Walter, J. F., III. (2022). Artificial attraction: Linking vessel monitoring system and habitat data to assess commercial exploitation on artificial structures in the Gulf of Mexico. *Frontiers in Marine Science*, *9*, Article 772292. https://doi.org/10.3389/fmars.2022.772292

Gill, A. B., Degraer, S., Lipsky, A., Mavraki, N., Methratta, E., & Brabant, R. (2020). Setting the context for offshore wind development on fish and fisheries. *Oceanography*, *33*, 118–127. https://doi.org/10.5670/oceanog.2020.411

Gill, D. A., Ritchie, L. A., Picou, J. S., Langhinrichsen-Rohling, J., Long, M. A., & Shenesey, J. W. (2014). The Exxon and BP oil spills: A comparison of psychosocial impacts. *Natural Hazards*, *74*, 1911–1932. https://doi.org/10.1007/s11069-014-1280-7

Glarou, M., Zrust, M., & Svendsen, J. C. (2020). Using artificial-reef knowledge to enhance the ecological function of offshore wind turbine foundations: Implications for fish abundance and diversity. *Journal of Marine Science and Engineering*, *8*, Article 332. https://doi.org/10.3390/jmse8050332

Glenn, H. D., Cowan, J. H., & Powers, J. E. (2017). A comparison of Red Snapper reproductive potential in the northwestern Gulf of Mexico: Natural versus artificial habitats. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science*, *9*, 139–148. https://doi.org/10.1080/19425120.2017.1282896

Graham, W. M., Martin, D. L., Felder, D. L., Asper, V. L., & Perry, H. M. (2003). Ecological and economic implications of a tropical jellyfish invader in the Gulf of Mexico. *Biological Invasions*, *5*, 53–69. https://doi.org/10.1023/A:1024046707234

Gray, T., Haggett, C., & Bell, D. (2005). Offshore windfarms and commercial fisheries in the UK: A study in stakeholder consultation. *Ethics, Place, and Environment*, *8*, 127–140. https://doi.org/10.1080/13668790500237013

Gray, M., Stromberg, P. L., & Rodmell, D. (2016). *Changes to fishing practices around the UK as a result of the development of offshore windfarms—Phase 1 (revised)*. The Crown Estate.

Grüss, A., Perryman, H. A., Babcock, E. A., Sagarese, S. R., Thorson, J. T., Ainsworth, C. H., Anderson, E. J., Brennan, K., Campbell, M. D., Christman, M. C., & Cross, S. (2018). Monitoring programs of the US Gulf of Mexico: Inventory, development and use of a large monitoring database to map fish and invertebrate spatial distributions. *Reviews in Fish Biology and Fisheries*, *28*, 667–691. https://doi.org/10.1007/s11160-018-9525-2

Gulf of Mexico Fishery Management Council. (2023). *Update to Red Snapper calibrations and Gray Snapper catch limits. Final framework action under the Fishery Management Plan for reef fish resources of the Gulf of Mexico*. https://www.fisheries.noaa.gov/s3//2024-05/Red-gray-snapper-12-20-23-final-wRevRFA.pdf

Gulf of Mexico Fishery Management Council. (2024). *Gulf of Mexico Fishery Management Council full council minutes. April 2024 meeting, Gulf Shores, Alabama, USA*. https://gulfcouncil.org/wp-content/uploads/GMFMC-Full-Council-April-2024_CMS.pdf

Gustafson, P. (2001). Meanings of place: Everyday experience and theoretical conceptualizations. *Journal of Environmental Psychology*, *21*, 5–16. https://doi.org/10.1006/jevp.2000.0185

Haas, H. L., Rose, K. A., Fry, B., Minello, T. J., & Rozas, L. P. (2004). Brown shrimp on the edge: Linking habitat to survival using an individual-based simulation model. *Ecological Applications*, *14*, 1232–1247. https://doi.org/10.1890/03-5101

Haase, S., von Dorrien, C., Kaljuste, O., Plantener, N., Sepp, E., Stelzenmüller, V., Velasco, A., & Oesterwind, D. (2023). The rapid expansion of offshore wind farms challenges the reliability of ICES-coordinated fish surveys—Insights from the Baltic Sea. *ICES Journal of Marine Science*, *82*, Article fsad124. https://doi.org/10.1093/icesjms/fsad124

Hanisko, D. S., Steen, M., Kellison, G. T., Walter, J. F., III, Segarra, K., Logan, M., Smith, A., Blyth, B., Howson, U., Paxton, A. B., Silverman, N., & Rosegger, B. (2025). *NOAA Fisheries and BOEM survey mitigation strategy: Southeast U.S. region* [In internal review]. National Oceanic and Atmospheric Administration.

Harris, H. E., Patterson, W. F., III, Ahrens, R. N., & Allen, M. S. (2019). Detection and removal efficiency of invasive lionfish in the northern Gulf of Mexico. *Fisheries Research*, *213*, 22–32. https://doi.org/10.1016/j.fishres.2019.01.002

Harris, H. E., Patterson, W. F., III, Ahrens, R. N., Allen, M. S., Chagaris, D. D., & Larkin, S. L. (2023). The bioeconomic paradox of market-based invasive species harvest: A case study of the commercial lionfish fishery. *Biological Invasions*, *25*, 1595–1612. https://doi.org/10.1007/s10530-023-02998-5

Harsanyi, P., Scott, K., Easton, B. A., de la Cruz Ortiz, G., Chapman, E. C., Piper, A. J., Rochas, C. M., & Lyndon, A. R. (2022). The effects of anthropogenic electromagnetic fields (EMF) on the early development of two commercially important crustaceans, European lobster, *Homarus gammarus (L.)* and edible crab, *Cancer pagurus (L.)*. *Journal of Marine Science and Engineering*, *10*, Article 564. https://doi.org/10.3390/jmse10050564

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Hawkins, A. D., Roberts, L., & Cheesman, S. (2014). Responses of free-living coastal pelagic fish to impulsive sounds. *The Journal of the Acoustical Society of America*, *135*, 3101–3116. https://doi.org/10.1121/1.4870697

Hazen, E. L., Craig, J. K., Good, C. P., & Crowder, L. B. (2009). Vertical distribution of fish biomass in hypoxic waters on the Gulf of Mexico shelf. *Marine Ecology Progress Series*, *375*, 195–207. https://doi.org/10.3354/meps07791

Hendon, R., Serafy, J., Walter, J., Lipsky, A., Curtis, T., Di Natale, A., Rouyer, T., Hanke, A., Alvarez-Berastegui, D., Orbesen, E., Lauretta, M., & Steizenmüller, V. (2024). Offshore wind energy development and highly migratory species: Ecological fishery, and management implications. *ICCAT Collective Volume of Scientific Papers*, *81*, 1–21.

Hermans, A., Winter, H. V., Gill, A. B., & Murk, A. J. (2024). Do electromagnetic fields from subsea power cables affect benthic elasmobranch behaviour? A risk-based approach for the Dutch continental shelf. *Environmental Pollution*, *346*, Article 123570. https://doi.org/10.1016/j.envpol.2024.123570

Hernandez, F. J., Shaw, R. F., Cope, J. S., Ditty, J. G., Farooqi, T., & Benfield, M. C. (2003). The across-shelf larval, postlarval, and juvenile fish assemblages collected at offshore oil and gas platforms west of the Mississippi river delta. In D. Stanley, & A. Scarborough-Bull (Eds.), *Fisheries, reefs, and offshore development* (Symposium 36, pp. 39–72). American Fisheries Society.

Heyman, W. D., Grüss, A., Biggs, C. R., Kobara, S., Farmer, N. A., Karnauskas, M., Lowerre-Barbieri, S., & Erisman, B. (2019). Cooperative monitoring, assessment, and management of fish spawning aggregations and associated fisheries in the U.S. Gulf of Mexico. *Marine Policy*, *109*, Article 103689. https://doi.org/10.1016/j.marpol.2019.103689

Hicks, W. (2024, July 11). *Offshore wind turbines offer path for clean hydrogen production*. National Renewable Energy Laboratory. https://www.nrel.gov/news/detail/program/2024/offshore-wind-turbines-offer-path-for-clean-hydrogen-production

Hobbs, B. F., Ludsin, S. A., Knight, R. L., Ryan, P. A., Biberhofer, J., & Ciborowski, J. J. (2002). Fuzzy cognitive mapping as a tool to define management objectives for complex ecosystems. *Ecological Applications*, *12*, 1548–1565. https://doi.org/10.1890/1051-0761(2002)012[1548:FCMAAT]2.0.CO;2

Hogan, F., Hooker, B., Jensen, B., Johnston, L., Lipsky, A., Methratta, E., Silva, A., & Hawkins, A. (2023). *Fisheries and offshore wind interactions: Synthesis of science* (Technical Memorandum NMFS-NE 291). National Oceanic and Atmospheric Administration. https://doi.org/10.25923/tcjt-3a69

Hooper, T., Hattam, C., & Austen, M. (2017). Recreational use of offshore wind farms: Experiences and opinions of sea anglers in the UK. *Marine Policy*, *78*, 55–60. https://doi.org/10.1016/j.marpol.2017.01.013

Hutchison, Z. L., Bartley, M. L., Degraer, S., English, P., Khan, A., Livermore, J., Rumes, B., & King, J. W. (2020b). Offshore wind energy and benthic habitat changes. *Oceanography*, *33*, 58–69. https://doi.org/10.5670/oceanog.2020.406

Hutchison, Z. L., Gill, A. B., Sigray, P., He, H., & King, J. W. (2020c). Anthropogenic electromagnetic fields (EMF) influence the behaviour of bottom-dwelling marine species. *Scientific Reports*, *10*, Article 4219. https://doi.org/10.1038/s41598-020-60793-x

Hutchison, Z. L., Secor, D. H., & Gill, A. B. (2020a). The interaction between resource species and electromagnetic fields associated with electricity production by offshore wind farms. *Oceanography*, *33*, 96–107. https://doi.org/10.5670/oceanog.2020.409

Ivanov, E., Capet, A., De Borger, E., Degraer, S., Delhez, E. J., Soetaert, K., Vanaverbeke, J., & Grégoire, M. (2021). Offshore wind farm footprint on organic and mineral particle flux to the bottom. *Frontiers in Marine Science*, *8*, Article 631799. https://doi.org/10.3389/fmars.2021.631799

Janßen, H., Schröder, T., Zettler, M. L., & Pollehne, F. (2015). Offshore wind farms in the southwestern Baltic Sea: A model study of regional impacts on oxygen conditions. *Journal of Sea Research*, *95*, 248–257. https://doi.org/10.1016/j.seares.2014.05.001

Johnston, M. W., Bernard, A. M., & Shivji, M. S. (2017). Forecasting lionfish sources and sinks in the Atlantic: Are Gulf of Mexico reef fisheries at risk? *Coral Reefs*, *36*, 169–181. https://doi.org/10.1007/s00338-016-1511-3

Jorgensen, B. S., & Stedman, R. C. (2001). Sense of place as an attitude: Lakeshore owners attitudes toward their properties. *Journal of Environmental Psychology*, *21*, 233–248. https://doi.org/10.1006/jevp.2001.0226

Karnauskas, M., Kelble, C. R., Regan, S., Quenée, C., Allee, R., Jepson, M., Freitag, A., Craig, J. K., Carollo, C., & Barbero, L. (2017). *2017 ecosystem status report update for the Gulf of Mexico* (Technical Memorandum NMFS-SEFSC-706). National Oceanic and Atmospheric Administration. https://www.aoml.noaa.gov/ocd/ocdweb/ESR_GOMIEA/report/GoMEcosystemStatusReport2017_NMFS-SEFSC-706_FINAL.pdf

Karnauskas, M., Schirripa, M. J., Craig, J. K., Cook, G. S., Kelble, C. R., Agar, J. J., Black, B. A., Enfield, D. B., Lindo-Atichati, D., Muhling, B. A., & Purcell, K. M. (2015). Evidence of climate-driven ecosystem reorganization in the Gulf of Mexico. *Global Change Biology*, *21*, 2554–2568. https://doi.org/10.1111/gcb.12894

Karnauskas, M., Schirripa, M. J., Kelble, C. R., Cook, G. S., & Craig, K. J. (2013). *Ecosystem status report for the Gulf of Mexico* (Technical Memorandum NMFS-SEFSC-653). National Oceanic and Atmospheric Administration.

Karnauskas, M., Walter, J. F., III, Campbell, M. D., Pollack, A. G., Drymon, J. M., & Powers, S. (2017). Red Snapper distribution on natural habitats and artificial structures in the northern Gulf of Mexico. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science*, *9*, 50–67. https://doi.org/10.1080/19425120.2016.1255684

Keller, B. A., Putman, N. F., Grubbs, R. D., Portnoy, D. S., & Murphy, T. P. (2021). Map-like use of earth's magnetic field in sharks. *Current Biology*, *31*, 2881–2886. https://doi.org/10.1016/j.cub.2021.03.103

Kerckhof, F., Rumes, B., & Degraer, S. (2019). About "mytilisation" and "slimeification": A decade of succession of the fouling assemblages on wind turbines off the Belgian coast. In S. Degraer, R. Brabant, B. Rumes, & L. Vigin (Eds.), *Environmental impacts of offshore wind farms in the Belgian part of the North Sea: Marking a decade of monitoring, research and innovation* (pp. 73–84). Royal Belgian Institute of Natural Sciences.

Kerckhof, F., Rumes, B., Jacques, T., Degraer, S., & Noro, A. (2010). Early development of the subtidal marine biofouling on a concrete offshore windmill foundation on the Thornton Bank (southern North Sea): First monitoring results. *International Journal of the Society for Underwater Technology*, *29*, 137–149. https://doi.org/10.3723/ut.29.137

Kerckhof, F., Rumes, B., Norro, A., Houziaux, J.-S., & Degraer, S. (2012). A comparison of the first stages of biofouling in two offshore wind farms in the Belgian part of the North Sea. In S. Degraer (Ed.), *Offshore wind farms in the Belgian part of the North Sea: Heading for an understanding of environmental impacts* (pp. 17–39). Royal Belgian Institute for Natural Sciences.

Kilborn, J. P. (2024). Characterizing a regional fishery ecosystem trajectory: The Humpty Dumpty fish tale of the U.S. Gulf of Mexico from 1986 to 2013. *Frontiers in Marine Science*, *11*, Article 1333756. https://doi.org/10.3389/fmars.2024.1333756

Kilborn, J. P., Drexler, M., & Jones, D. L. (2018). Fluctuating fishing intensities and climate dynamics reorganize the Gulf of Mexico's fisheries resources. *Ecosphere*, *9*, Article e02487. https://doi.org/10.1002/ecs2.2487

Kiørboe, T., & MacKenzie, B. (1995). Turbulence-enhanced prey encounter rates in larval fish: Effects of spatial scale, larval behaviour and size. *Journal of Plankton Research*, *17*, 2319–2331. https://doi.org/10.1093/plankt/17.12.2319

Kirkpatrick, A. J., Benjamin, S., DePiper, G. S., Murphy, T., Steinback, S., & Demarest, C. (2017). *Socioeconomic impact of outer continental shelf wind energy development on fisheries in the U.S. Atlantic. Volume I—Report narrative* (OCS Study BOEM 2017-012). Bureau of Ocean Energy Management.

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Klain, S. C., Beveridge, R., & Bennett, N. J. (2014). Ecologically sustainable but unjust? Negotiating equity and authority in common-pool marine resource management. *Ecology and Society*, *19*, Article 52. https://doi.org/10.5751/ES-07123-190452

Klajbor, W., Harris, H. E., Freitag, A., Troast, B., Turley, B., & Kelble, C. (2025). Winds of change?: A stakeholder-driven conceptual model of offshore wind energy development in the northwestern Gulf of Mexico [Manuscript in preparation].

Krone, R., Dederer, G., Kanstinger, P., Krämer, P., Schneider, C., & Schmalenbach, I. (2017). Mobile demersal megafauna at common offshore wind turbine foundations in the German bight (North Sea) two years after deployment-increased production rate of *Cancer pagurus*. *Marine Environmental Research*, *123*, 53–61. https://doi.org/10.1016/j.marenvres.2016.11.011

Lages, B. G., Fleury, B. G., Menegola, C., & Creed, J. C. (2011). Change in tropical rocky shore communities due to an alien coral invasion. *Marine Ecology Progress Series*, *438*, 85–96. https://doi.org/10.3354/meps09290

Layman, C. A., & Allgeier, J. E. (2020). An ecosystem ecology perspective on artificial reef production. *Journal of Applied Ecology*, *57*, 2139–2148. https://doi.org/10.1111/1365-2664.13748

Layman, C. A., Allgeier, J. E., & Montaña, C. G. (2016). Mechanistic evidence of enhanced production on artificial reefs: A case study in a Bahamian seagrass ecosystem. *Ecological Engineering*, *95*, 574–579. https://doi.org/10.1016/j.ecoleng.2016.06.109

Lee, M. R., & Blanchard, T. C. (2012). Community attachment and negative affective states in the context of the BP deepwater horizon disaster. *The American Behavioral Scientist*, *56*, 24–47. https://doi.org/10.1177/0002764211409384

Lefaible, N., Braeckman, U., Degraer, S., Vanaverbeke, J., & Moens, T. (2023). A wind of change for soft-sediment infauna within operational offshore windfarms. *Marine Environmental Research*, *188*, Article 106009. https://doi.org/10.1016/j.marenvres.2023.106009

Leiva, L., Scholz, S., Giménez, L., Boersma, M., Torres, G., Krone, R., & Tremblay, N. (2021). Noisy waters can influence young-of-year lobsters' substrate choice and their antipredatory responses. *Environmental Pollution*, *291*, Article 118108. https://doi.org/10.1016/j.envpol.2021.118108

Lemasson, A. J., Somerfield, P. J., Schratzberger, M., Thompson, M. S. A., Firth, L. B., Couce, E., McNeill, C. L., Nunes, J., Pascoe, C., Watson, S. C. L., & Knights, A. M. (2024). A global meta-analysis of ecological effects from offshore marine artificial structures. *Nature Sustainability*, *7*, 485–495. https://doi.org/10.1038/s41893-024-01311-z

Li, O., Gray, S. A., & Sutton, S. G. (2016). Mapping recreational fishers' informal learning of scientific information using a fuzzy cognitive mapping approach to mental modelling. *Fisheries Management and Ecology*, *23*, 315–329. https://doi.org/10.1111/fme.12174

Lindquist, D. C., Shaw, R. F., & Hernandez, F. J., Jr. (2005). Distribution patterns of larval and juvenile fishes at offshore petroleum platforms in the north-central Gulf of Mexico. *Estuarine, Coastal and Shelf Science*, *62*, 655–665. https://doi.org/10.1016/j.ecss.2004.10.001

Louisiana Department of Wildlife and Fisheries. (2020). *Update assessment of Striped Mullet (*Mugil cephalus*) in Louisiana Waters*. https://www.wlf.louisiana.gov/assets/Resources/Publications/Stock_Assesments/Striped_Mullet/2020_Update_Assessment_of_Striped_Mullet.pdf

Lowerre-Barbieri, S. K., Friess, C., Griffin, L. P., Morley, D., Skomal, G. B., Bickford, J. W., Hammerschlag, N., Rider, M. J., Smukall, M. J., van Zinnicq Bergmann, M. P. M., Guttridge, T. L., Kroetz, A. M., Grubbs, R. D., Gervasi, C. L., Rehage, J. S., Poulakis, G. R., Bassos-Hull, K., Gardiner, J. M., Casselberry, G. A., … Brownscombe, J. W. (2021). Movescapes and eco-evolutionary movement strategies in marine fish: Assessing a connectivity hotspot. *Fish and Fisheries*, *22*, 1321–1344. https://doi.org/10.1111/faf.12589

Lu, Z., Li, G., Liu, Z., & Wang, L. (2022). Offshore wind farms changed the spatial distribution of chlorophyll-a on the sea surface. *Frontiers in Marine Science*, *9*, Article 1008005. https://doi.org/10.3389/fmars.2022.1008005

Mackinson, S., Curtis, H., Brown, R., McTaggart, K., Taylor, N., Neville, S., & Rogers, S. (2006). *A report on the perceptions of the fishing industry into the potential socio-economic impacts of offshore wind energy developments on their work patterns and income* (Science Series Technical Report 133). Cefas. https://www.cefas.co.uk/publications/techrep/tech133.pdf

MacNeil, M. A., Chapman, D. D., Heupel, M., Simpfendorfer C. A., Heithaus M., Meekan M., Harvey E., Goetze J., Kiszka J., Bond M. E., Currey-Randall L. M., Speed C. W., Sherman C. S., Rees M. J., Udyawer V., Flowers K. I., Clementi G., Valentin-Albanese J., Gorham T. … Cinner J. E. (2020). Global status and conservation potential of reef sharks. *Nature*, *583*, 801–806. https://doi.org/10.1038/s41586-020-2519-y

Magnuson–Stevens Fishery Conservation and Management Act. (1976). 16 U.S.C. § 1801 *et seq*.

Mareska, J., & Bellais, D. (2023). *Wenchman: Gulf of Mexico. Presentation to the Gulf of Mexico Fishery Management Council's Scientific and Statistical Committee, March 2023, Tampa, Florida, USA*. Gulf of Mexico Fishery Management Council. https://gulfcouncil.org/wp-content/uploads/B-4e-Wenchman-fishery-and-trawl-landings.pdf

Marshall, A., Barreto, R., Carlson, J., Fernando, D., Fordham, S., Francis, M. P., Derrick, D., Herman, K., Jabado, R. W., Liu, K. M., Rigby, C. L. & Romanov, E. (2022). Mobula birostris (amended version of 2020 assessment). *The IUCN Red List of Threatened Species*, *2022*, Article e.T198921A214397182. https://dx.doi.org/10.2305/IUCN.UK.2022-1.RLTS.T198921A214397182.en

McEachran, J. D., & Fechhelm, F. D. (1998). *Fishes of the Gulf of Mexico* (Vol. 1). University of Texas Press.

McEachran, J. D., & Fechhelm, F. D. (2006). *Fishes of the Gulf of Mexico* (Vol. 2). University of Texas Press.

McLean, D. L., Ferreira, L. C., Benthuysen, J. A., Miller, K. J., Schläppy, M. L., Ajemian, M. J., Berry, O., Birchenough, S. N., Bond, T., Boschetti, F., & Bull, A. S. (2022). Influence of offshore oil and gas structures on seascape ecological connectivity. *Global Change Biology*, *28*, 3515–3536. https://doi.org/10.1111/gcb.16134

Methratta, E. T., & Dardick, W. R. (2019). Meta-analysis of finfish abundance at offshore wind farms. *Reviews in Fisheries Science & Aquaculture*, *27*, 242–260. https://doi.org/10.1080/23308249.2019.1584601

Methratta, E. T., Hawkins, A., Hooker, B. R., Lipsky, A., & Hare, J. A. (2020). Offshore wind development in the northeast US shelf large marine ecosystem. *Oceanography*, *33*, 16–27. https://doi.org/10.5670/oceanog.2020.402

Methratta, E. T., Silva, A., Lipsky, A., Ford, K., Christel, D., & Pfeiffer, L. (2023). Science priorities for offshore wind and fisheries research in the Northeast U.S. Continental Shelf ecosystem: Perspectives from scientists at the National Marine Fisheries Service. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science*, *15*, Article e10242. https://doi.org/10.1002/mcf2.10242

Meyer, G. H., & Franks, J. S. (1996). Food of Cobia, *Rachycentron canadum*, from the northcentral Gulf of Mexico. *Gulf and Caribbean Research*, *9*, 161–167. https://doi.org/10.18785/grr.0903.02

Montero, J. T., Chesney, T. A., Bauer, J. R., Froeschke, J. T., & Graham, J. (2016). Brown shrimp (*Farfantepenaeus aztecus*) density distribution in the northern Gulf of Mexico: An approach using boosted regression trees. *Fisheries Oceanography*, *25*, 337–348. https://doi.org/10.1111/fog.12156

Muller-Karger, F. E., Smith, J. P., Werner, S., Chen, R., Roffer, M., Liu, Y., Muhling, B., Lindo-Atichati, D., Lamkin, J., Cerdeira-Estrada, S., & Enfield, D. B. (2015). Natural variability of surface oceanographic conditions in the offshore Gulf of Mexico. *Progress in Oceanography*, *134*, 54–76. https://doi.org/10.1016/j.pocean.2014.12.007

Musial, W., Spitsen, P., Duffy, P., Beiter, P., Shields, M., Hernando, D. M., Hammond, R., Marquis, M., King, J., & Sathish, S. (2023). *Offshore wind market report: 2023 edition*. Department of Energy, Office of Energy Efficiency and Renewable Energy. https://www.energy.gov/sites/default/files/2023-09/doe-offshore-wind-market-report-2023-edition.pdf

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Nagel, T., Chauchat, J., Wirth, A., & Bonamy, C. (2018). On the multi-scale interactions between an offshore-wind-turbine wake and the ocean-sediment dynamics in an idealized framework–a numerical investigation. *Renewable Energy*, *115*, 783–796. https://doi.org/10.1016/j.renene.2017.08.078

National Marine Fisheries Service. (2023). *Fisheries economics of the United States 2020* (Technical Memorandum NMFS-F/SPO-236A). National Oceanic and Atmospheric Administration. https://media.fisheries.noaa.gov/2024-01/FEUS-2020-final2-web.pdf

National Marine Fisheries Service. (2024a). *Fisheries of the United States, 2022* (Current Fishery Statistics No. 2022). https://www.fisheries.noaa.gov/national/sustainable-fisheries/fisheries-united-states

National Marine Fisheries Service. (2024b). *Status of stocks 2023: Annual Report to Congress on the status of U.S. fisheries*. https://www.fisheries.noaa.gov/s3/2024-04/2023SOS-final.pdf

National Marine Fisheries Service Fisheries Office of Science and Technology. *Commercial landings query*. Landings [Online database]. Retrieved March, 2024, from www.fisheries.noaa.gov/foss

National Oceanic and Atmospheric Administration. (2023). *Coastal population*. National Marine Ecosystem Status website. https://ecowatch.noaa.gov/thematic/coastal-population

Nishimoto, M. M., Simons, R. D., & Love, M. S. (2019). Offshore oil production platforms as potential sources of larvae to coastal shelf regions off southern California. *Bulletin of Marine Science*, *95*, 535–558. https://doi.org/10.5343/bms.2019.0033

Office for Coastal Management. (2024). *Offshore oil and gas platforms* [Online data set]. National Oceanic and Atmospheric Administration's National Centers for Environmental Information. Retrieved September, 2024, from https://www.fisheries.noaa.gov/inport/item/66162

Osborn, K. W., Maghan, B. W., & Drummond, S. B. (1969). *Gulf of Mexico shrimp atlas* (Circular 312). U.S. Department of the Interior, Bureau of Commercial Fisheries.

Osenberg, C. W., St. Mary, C. M., Wilson, J. A., & Lindberg, W. J. (2002). A quantitative framework to evaluate the attraction–production controversy. *ICES Journal of Marine Science*, *59*, S214–S221. https://doi.org/10.1006/jmsc.2002.1222

Osowski, A. R., & Szedlmayer, S. T. (2022). Red Snapper (*Lutjanus campechanus*) abundance on oil and gas platforms based on mark–recapture methods in the northern Gulf of Mexico. *Canadian Journal of Fisheries and Aquatic Sciences*, *79*, 1546–1560. https://doi.org/10.1139/cjfas-2021-0227

Ørsted. (2019). *Making green energy affordable: How the offshore wind energy industry matured—and what we can learn from it* [White paper]. Ørsted. https://orsted.com/en/insights/white-papers/making-green-energy-affordable

Outer Continental Shelf Lands Act. (1953). 43 U.S.C. § 1331 *et seq*.

Oxenford, H. A., & Hunte, W. (1999). Feeding habits of the Dolphinfish (*Coryphaena hippurus*) in the eastern Caribbean. *Scientia Marina*, *63*, 303–315. https://doi.org/10.3989/scimar.1999.63n3-4317

Patterson, W. F., III. (2007). A review of movement in Gulf of Mexico Red Snapper: Implications for population structure. In W. F. Patterson III, J. H. Cowan Jr., G. R. Fitzhugh, & D. L. Nieland (Eds.), *Red snapper ecology and fisheries in the U.S. Gulf of Mexico* (Symposium 60, pp. 221–235). American Fisheries Society. https://doi.org/10.47886/9781888569971.ch16

Patterson, W. F., Robinson, K. L., Barnett, B. K., Campbell, M. D., Chagaris, D. C., Chanton, J. P., Daly, K. L., Hanisko, D. S., Hernandez, F. J., Murawski, S. A., Pollack, A. G., Portnoy, D. S., & Pulster, E. L. (2023). Evidence of population-level impacts and resiliency for Gulf of Mexico shelf taxa following the deepwater horizon oil spill. *Frontiers in Marine Science*, *10*, Article 1198163. https://doi.org/10.3389/fmars.2023.1198163

Pattillo, M. E., Czapla, T. E., Nelson, D. M., & Monaco, M. E. (1997). *Distribution and abundance of fishes and invertebrates in Gulf of Mexico estuaries. Volume 2: Species life history summaries*. National Oceanic and Atmospheric Administration, National Ocean Service.

Paxton, A. B., Peterson, C. H., Taylor, J. C., Adler, A. M., Pickering, E. A., & Silliman, B. R. (2019). Artificial reefs facilitate tropical fish at their range edge. *Communications Biology*, *2*, Article 168. https://doi.org/10.1038/s42003-019-0398-2

Paxton, A. B., III, Runde, B. J., Smith, C. S., Lester, S. E., Vozzo, M. L., Saunders, M. I., Steward, D. N., Lemoine, H. R., Valdez, S. R., Gittman, R. K., Narayan, S., Allgeier, J., Morris, R. L., Nowacek, D. P., Seaman, W., Halpin, P. N., Angelini, C., & Silliman, B. R. (2025). Leveraging built marine structures to benefit and minimize impacts on natural habitats. *BioScience*, *75*, 172–183. https://doi.org/10.1093/biosci/biae135

Paxton, A. B., Steward, D. a. N., Mille, K. J., Renchen, J., Harrison, Z. H., Byrum, J. S., Brinton, C., Nelson, A., Simpson, E., Clarke, P. J., LaPorta, C., Barrett, P. D., Rousseau, M., Newton, D. C., Rigby, R. B., Williams, D. T., Shipley, J. B., Murakawa, P., Runde, B. J., … Taylor, J. C. (2024). Artificial reef footprint in the United States ocean. *Nature Sustainability*, *7*, 140–147. https://doi.org/10.1038/s41893-023-01258-7

Perry, R. M., & Heyman, W. D. (2020). Considerations for offshore wind development effects on fisheries in the United States: A review of existing studies, new efforts, and opportunities for innovation. *Oceanography*, *33*, 28–37. https://doi.org/10.5670/oceanog.2020.403

Petersen, J. K., & Malm, T. (2006). Offshore windmill farms: Threats to or possibilities for the marine environment. *AMBIO: A Journal of the Human Environment*, *35*, 75–80. https://doi.org/10.1579/0044-7447(2006)35[75:OWFTTO]2.0.CO;2

Pickering, H., & Whitmarsh, D. (1997). Artificial reefs and fisheries exploitation: A review of the 'attraction versus production' debate, the influence of design and its significance for policy. *Fisheries Research*, *31*, 39–59. https://doi.org/10.1016/S0165-7836(97)00019-2

Pierce, S. J., & Norman, B. (2016). Rhincodon typus. *The IUCN Red List of Threatened Species*, *2016*, Article e.T19488A2365291. https://dx.doi.org/10.2305/IUCN.UK.2016-1.RLTS.T19488A2365291.en

Pinsky, M. L., Fenichel, E., Fogarty, M., Levin, S., McCay, B., St. Martin, K., Selden, R. L., & Young, T. (2021). Fish and fisheries in hot water: What is happening and how do we adapt? *Population Ecology*, *63*, 17–26. https://doi.org/10.1002/1438-390X.12050

Pinsky, M. L., Worm, B., Fogarty, M. J., Sarmiento, J. L., & Levin, S. A. (2013). Marine taxa track local climate velocities. *Science*, *341*, 1239–1242. https://doi.org/10.1126/science

Popper, A. N., Hice-Dunton, L., Jenkins, E., Higgs, D. M., Krebs, J., Mooney, A., Rice, A., Roberts, L., Thomsen, F., Vigness-Raposa, K., Zeddies, D., & Williams, K. A. (2022). Offshore wind energy development: Research priorities for sound and vibration effects on fishes and aquatic invertebrates. *The Journal of the Acoustical Society of America*, *151*, 205–215. https://doi.org/10.1121/10.0009237

Portnoy, D. S., Fields, A. T., Puritz, J. B., Hollenbeck, C. M., & Patterson, W. F., III. (2022). Genomic analysis of Red Snapper, *Lutjanus campechanus*, population structure in the U.S. Atlantic and Gulf of Mexico. *ICES Journal of Marine Science*, *79*, 12–21. https://doi.org/10.1093/icesjms/fsab239

Powers, S. P., Drymon, J. M., Hightower, C. L., Spearman, T., Bosarge, G. S., & Jefferson, A. (2018). Distribution and age composition of Red Snapper across the inner continental shelf of the north-central Gulf of Mexico. *Transactions of the American Fisheries Society*, *147*, 791–805. https://doi.org/10.1002/tafs.10081

Powers, S. P., Peterson, C. H., Cebrian, J., & Heck, K. L., Jr. (2017). Response of nearshore ecosystems to the Deepwater Horizon oil spill. *Marine Ecology Progress Series*, *576*, 107–110. https://doi.org/10.3354/meps12254

Püts, M., Kempf, A., Möllmann, C., & Taylor, M. (2023). Trade-offs between fisheries, offshore wind farms and marine protected areas in the southern North Sea – Winners, losers and effective spatial management. *Marine Policy*, *152*, Article 105574. https://doi.org/10.1016/j.marpol.2023.105574

Rabalais, N. N., Turner, R. E., & Wiseman, W. J., Jr. (2001). Hypoxia in the Gulf of Mexico. *Journal of Environmental Quality*, *30*, 320–329. https://doi.org/10.2134/jeq2001.302320x

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

Randall, A. L., Jossart, J. A., Matthews, T., Steen, M., Boube, I., Stradley, S., Del Rio, R., Inzinna, D., Oos, C., Coats, L., Shin, G., Griffith, C., & Morris, J. A., Jr. (2022). *A wind energy area siting analysis for the Gulf of Mexico call area*. https://www.boem.gov/sites/default/files/documents/renewable-energy/state-activities/GOM-WEA-Modeling-Report-Combined.pdf

Raoux, A., Tecchio, S., Pezy, J. P., Lassalle, G., Degraer, S., Wilhelmsson, D., Cachera, M., Ernande, B., Le Guen, C., Haraldsson, M., & Grangeré, K. (2017). Benthic and fish aggregation inside an offshore wind farm: Which effects on the trophic web functioning? *Ecological Indicators*, *72*, 33–46. https://doi.org/10.1016/j.ecolind.2016.07.037

Reeves, D. B., Chesney, E. J., Munnelly, R. T., Baltz, D. M., & Marx, B. D. (2018). Abundance and distribution of reef-associated fishes around small oil and gas platforms in the northern Gulf of Mexico's hypoxic zone. *Estuaries and Coasts*, *41*, 1835–1847. https://doi.org/10.1007/s12237-017-0349-4

Reubens, J. T., Degraer, S., & Vincx, M. (2014). The ecology of benthopelagic fishes at offshore wind farms: A synthesis of 4 years of research. *Hydrobiologia*, *727*, 121–136. https://doi.org/10.1007/s10750-013-1793-1

Reubens, J. T., Pasotti, F., Degraer, S., & Vincx, M. (2013). Residency, site fidelity and habitat use of Atlantic Cod (*Gadus morhua*) at an offshore wind farm using acoustic telemetry. *Marine Environmental Research*, *90*, 128–135. https://doi.org/10.1016/j.marenvres.2013.07.001

Rigby, C. L., Barreto, R., Carlson, J., Fernando, D., Fordham, S., Francis, M. P., Herman, K., Jabado, R. W., Liu, K. M., Marshall, A., Pacoureau, N., Romanov, E., Sherley, R. B., & Winker, H. (2019). Carcharhinus longimanus. *The IUCN Red List of Threatened Species*, *2019*, Article e.T39374A2911619. https://doi.org/10.2305/IUCN.UK.2019-3.RLTS.T39374A2911619.en

Roach, M., Revill, A., & Johnson, M. J. (2022). Co-existence in practice: A collaborative study of the effects of the westermost rough offshore wind development on the size distribution and catch rates of a commercially important lobster (*Homarus gammarus*) population. *ICES Journal of Marine Science*, *79*, 1175–1186. https://doi.org/10.1093/icesjms/fsac040

Röckmann, C., Ulrich, C., Dreyer, M., Bell, E., Borodzicz, E., Haapasaari, P., Hauge, K. H., Howell, D., Mäntyniemi, S., & Miller, D. (2012). The added value of participatory modelling in fisheries management – What has been learnt? *Marine Policy*, *36*, 1072–1085. https://doi.org/10.1016/j.marpol.2012.02.027

Rumes, B., Coates, D., De Mesel, I., Derweduwen, J., Kerckhof, F., Reubens, J., & Vandendriessche, S. (2013). Does it really matter? Changes in species richness and biomass at different spatial scales. In S. Degraer, R. Brabant, & B. Rumes (Eds.), *Environmental impacts of offshore wind farms in the Belgian part of the North Sea: Learning from the past to optimise future monitoring programmes* (pp. 183–189). Royal Belgian Institute of Natural Sciences.

Russell, A., Firestone, J., Bidwell, D., & Gardner, M. (2020). Place meaning and consistency with offshore wind: An island and coastal tale. *Renewable & Sustainable Energy Reviews*, *132*, Article 110044. https://doi.org/10.1016/j.rser.2020.110044

Sagarese, S. R., Nuttall, M. A., Geers, T. M., Lauretta, M. V., Walter, J. F., III, & Serafy, J. E. (2016). Quantifying the trophic importance of Gulf Menhaden within the northern Gulf of Mexico ecosystem. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science*, *8*, 23–45. https://doi.org/10.1080/19425120.2015.1091412

Samoteskul, K., Firestone, J., Corbett, J., & Callahan, J. (2014). Changing vessel routes could significantly reduce the cost of future offshore wind projects. *Journal of Environmental Management*, *141*, 146–154. https://doi.org/10.1016/j.jenvman.2014.03.026

Sandifer, P. A., Singer, B. H., & Colwell, R. R. (2021). The U.S. needs a national human health observing system. *Frontiers in Public Health*, *9*, Article 705597. https://doi.org/10.3389/fpubh.2021.705597

Scheld, A. M., Beckensteiner, J., Munroe, D. M., Powell, E. N., Borsetti, S., Hofmann, E. E., & Klinck, J. M. (2022). The Atlantic surfclam fishery and offshore wind energy development: 2. Assessing economic impacts. *ICES Journal of Marine Science*, *79*, 1801–1814. https://doi.org/10.1093/icesjms/fsac109

Schulze, A., Erdner, D. L., Grimes, C. J., Holstein, D. M., & Miglietta, M. P. (2020). Artificial reefs in the northern Gulf of Mexico: Community ecology amid the "ocean sprawl". *Frontiers in Marine Science*, *7*, Article 447. https://doi.org/10.3389/fmars.2020.00447

Schupp, M. F., Kafas, A., Buck, B. H., Krause, G., Onyango, V., Stelzenmüller, V., Davies, I., & Scott, B. E. (2021). Fishing within offshore wind farms in the north sea: Stakeholder perspectives for multi-use from Scotland and Germany. *Journal of Environmental Management*, *279*, Article 111762. https://doi.org/10.1016/j.jenvman.2020.111762

Scott, K., Harsanyi, P., & Lyndon, A. R. (2018). Understanding the effects of electromagnetic field emissions from marine renewable energy devices (MREDs) on the commercially important edible crab, *Cancer pagurus (L.)*. *Marine Pollution Bulletin*, *131*, 580–588. https://doi.org/10.1016/j.marpolbul.2018.04.062

SEDAR. (2020a). *SEDAR 28 update: Gulf of Mexico Cobia update stock assessment report*. https://sedarweb.org/documents/2019-update-sedar-28-gulf-of-mexico-cobia-final-report/

SEDAR. (2020b). *SEDAR 38 update: Gulf of Mexico King Mackerel update stock assessment report*. https://sedarweb.org/documents/2019-update-sedar-38-update-assessment-report-gulf-of-mexico-king-mackerel/

SEDAR. (2023). *SEDAR 81: Gulf of Mexico Spanish Mackerel operational assessment report. Southeast data, Assessment, and Review*. https://sedarweb.org/documents/sedar-81-gulf-of-mexico-spanish-mackerel-final-stock-assessment/

SEDAR. (2024). *SEDAR 85 stock assessment: Gulf of Mexico Yellowedge Grouper*.https://sedarweb.org/documents/sedar-85-gulf-of-mexico-yellowedge-grouper-final-stock-assessment-report/

Shaffer, M., Marriott, S., Lewis, K. A., Buszowski, J., & de Mutsert, K. (2023). Action science in practice: Co-production of a decision support tool visualizing effects of nutrient and hypoxia reduction goals on fisheries species. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science*, *15*, Article e10262. https://doi.org/10.1002/mcf2.10262

Shipp, R. L., & Bortone, S. A. (2009). A perspective of the importance of artificial habitat on the management of Red Snapper in the Gulf of Mexico. *Reviews in Fisheries Science*, *17*, 41–47. https://doi.org/10.1080/10641260802104244

Sinopoli, M., Battaglia, P., & Barreiros, J. P. (2017). Unusual presence of *Coryphaena hippurus* Linnaeus, 1758 (Perciformes: Coryphaenidae) under an offshore oil platform in southern Brazil. *Journal of Coastal Life Medicine*, *5*, 239–241. https://doi.org/10.12980/jclm.5.2017J7-48

Slotte, A., Hansen, K., Dalen, J., & Ona, E. (2004). Acoustic mapping of pelagic fish distribution and abundance in relation to a seismic shooting area off the Norwegian west coast. *Fisheries Research*, *67*, 143–150. https://doi.org/10.1016/j.fishres.2003.09.046

Smith, J. W., Hall, E. A., McNeill, N. A., & O'Bier, W. B. (2002). The distribution of purse-seine sets and catches in the Gulf Menhaden fishery in the northern Gulf of Mexico, 1994–98. *Gulf of Mexico Science*, *20*, Article 2. https://doi.org/10.18785/goms.2001.02

Smythe, T., Bidwell, D., & Tyler, G. (2021). Optimistic with reservations: The impacts of the United States' first offshore wind farm on the recreational fishing experience. *Marine Policy*, *127*, Article 104440. https://doi.org/10.1016/j.marpol.2021.104440

Snodgrass, D. J. G., Orbesen, E. S., Walter, J. F., III, Hoolihan, J. P., & Brown, C. A. (2020). Potential impacts of oil production platforms and their function as fish aggregating devices on the biology of highly migratory fish species. *Reviews in Fish Biology and Fisheries*, *30*, 405–422. https://doi.org/10.1007/s11160-020-09605-z

Soto, I. M., Cambazoglu, M. K., Boyette, A. D., Broussard, K., Sheehan, D., Howden, S. D., Shiller, A. M., Dzwonkowski, B., Hode, L., Fitzpatrick, P. J., & Arnone, R. A. (2018). Advection of *Karenia brevis* blooms from the Florida panhandle towards Mississippi coastal waters. *Harmful Algae*, *72*, 46–64. https://doi.org/10.1016/j.hal.2017.12.008

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

South Atlantic Fishery Management Council. (2024). *Establishing a fisheries fund regional administrator*. https://safmc.net/documents/fc2_a4a_special-initiative-on-offshore-wind-presentation_202406-pdf/

Special Initiative on Offshore Wind. (2024). *Fisheries mitigation project*. Retrieved September, 2024, from https://offshorewindpower.org/fisheries-mitigation-project

Stanley, D. R., & Wilson, C. A. (1989). Utilization of offshore platforms by recreational fishermen and scuba divers off the Louisiana coast. *Bulletin of Marine Science*, *44*, 767–776. https://www.ingentaconnect.com/contentone/umrsmas/bullmar/1989/00000044/00000002/art00020

Steenbeek, J., Romagnoni, G., Bentley, J., Heymans, J. J., Serpetti, N., Gonçalves, M., Santos, C., Warmelink, H., Mayer, I., Keijser, X., Fairgrieve, R., & Abspoel, L. (2020). Combining ecosystem modeling with serious gaming in support of transboundary maritime spatial planning. *Ecology and Society*, *25*, Article 21. https://doi.org/10.5751/ES-11580-250221

Stewart, J. D., Nuttall, M., Hickerson, E. L., & Johnston, M. A. (2018). Important juvenile Manta Ray habitat at Flower Garden Banks National Marine Sanctuary in the northwestern Gulf of Mexico. *Marine Biology*, *165*, Article 111. https://doi.org/10.1007/s00227-018-3364-5

Strayer, D. L., D'Antonio, C. M., Essl, F., Fowler, M. S., Geist, J., Hilt, S., Jarić, I., Jöhnk, K., Jones, C. G., Lambin, X., & Latzka, A. W. (2017). Boom-bust dynamics in biological invasions: Towards an improved application of the concept. *Ecology Letters*, *20*, 1337–1350. https://doi.org/10.1111/ele.12822

Streich, M. K., Ajemian, M. J., Wetz, J. J., Shively, J. D., Shipley, J. B., & Stunz, G. W. (2017). Effects of a new artificial reef complex on Red Snapper and the associated fish community: An evaluation using a before–after control–impact approach. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science*, *9*, 404–418. https://doi.org/10.1080/19425120.2017.1347116

Stunz, G. W., Patterson, W. F., III, Powers, S. P., Cowan, J. H., Jr., Rooker, J. R., Ahrens, R. A., Boswell, K., Carleton, L., Catalano, M., Drymon, J. M., Hoenig, J., Leaf, R., Lecours, V., Murawski, S., Portnoy, D., Saillant, E., Stokes, L. S., & Wells, R. J. D. (2021). *Estimating the absolute abundance of age-2+ Red Snapper (Lutjanus campechanus) in the U.S. Gulf of Mexico*. Mississippi-Alabama Sea Grant Consortium, NOAA Sea Grant. https://repository.library.noaa.gov/view/noaa/48520

Szedlmayer, S. T., & Shipp, R. L. (1994). Movement and growth of Red Snapper, *Lutjanus campechanus*, from an artificial reef area in the northeastern Gulf of Mexico. *Bulletin of Marine Science*, *55*, 887–896. https://www.ingentaconnect.com/content/umrsmas/bullmar/1994/00000055/f0020002/art00049

Taormina, B., Bald, J., Want, A., Thouzeau, G., Lejart, M., Desroy, N., & Carlier, A. (2018). A review of potential impacts of submarine power cables on the marine environment: Knowledge gaps, recommendations and future directions. *Renewable & Sustainable Energy Reviews*, *96*, 380–391. https://doi.org/10.1016/j.rser.2018.07.026

Tarnecki, J. H., Garner, S. B., & Patterson, W. F., III. (2021). Nonnative Regal Demoiselle, *Neopomacentrus cyanomos*, presence, abundance, and habitat factors in the north-central Gulf of Mexico. *Biological Invasions*, *23*, 1681–1693. https://doi.org/10.1007/s10530-020-02424-0

ten Brink, T. S., & Dalton, T. (2018). Perceptions of commercial and recreational fishers on the potential ecological impacts of the Block Island Wind Farm (US). *Frontiers in Marine Science*, *5*, Article 439. https://doi.org/10.3389/fmars.2018.00439

Tharp, R. M., Hostetter, N. J., Paxton, A. B., Taylor, J. C., & Buckel, J. A. (2024). Artificial structure selection by economically important reef fishes at North Carolina artificial reefs. *Frontiers in Marine Science*, *11*, Article 1373494. https://doi.org/10.3389/fmars.2024.1373494

The White House. (2021). Executive Order 14008 of January 27, 2021: Tackling the climate crisis at home and abroad. *Federal Register* 86:19(8 March 2021):7619–7633. https://www.regulations.gov/document/EPA-HQ-OPPT-2021-0202-0012

The White House. (2025). Temporary withdrawal of all areas on the outer continental shelf from offshore wind leasing and review of the federal government's leasing and permitting practices for wind projects. *Federal Register* 90:18(20 January 2025):8363–8365. https://www.govinfo.gov/content/pkg/FR-2025-01-29/pdf/2025-01966.pdf

Thorne, L. H., & Wiley, D. N. (2024). Evaluating drivers of recent large whale strandings on the East Coast of the United States. *Conservation Biology*, *38*, Article e14302. https://doi.org/10.1111/cobi.14302

Topping, D. T., & Szedlmayer, S. T. (2011). Site fidelity, residence time and movements of Red Snapper *Lutjanus campechanus* estimated with long-term acoustic monitoring. *Marine Ecology Progress Series*, *437*, 183–200. https://doi.org/10.3354/meps09293

Tricas, T. C., & Sisneros, J. A. (2004). Ecological functions and adaptations of the elasmobranch electrosense. In G. von der Emde, J. Mogdans, & B. J. Kapoor (Eds.), *The senses of fish* (pp. 308–329). Springer. https://doi.org/10.1007/978-94-007-1060-3_14

Tuan, Y.-F. (1974). *Topophilia: A study of environmental perception, attitudes, and values*. Prentice Hall.

Tunnell, J. W., Jr. (2017). Shellfish of the Gulf of Mexico. In C. H. Ward (Ed.), *Habitats and biota of the Gulf of Mexico: Before the Deepwater Horizon oil spill. Volume 1: Water quality, sediments, sediment contaminants, oil and gas seeps, coastal habitats, offshore plankton and benthos, and shellfish* (pp. 769–839). https://link.springer.com/chapter/10.1007/978-1-4939-3447-8_8

Tuttle, L. J. (2017). Direct and indirect effects of invasive lionfish on coral-reef cleaning mutualists. *Marine Ecology Progress Series*, *569*, 163–172. https://doi.org/10.3354/meps12092

U.S. Coast Guard. (2020). The areas offshore of MA and RI port access route study area USCG-2019-0131. *Federal Register* 85:102(27 May 2020):31792–31796. https://www.regulations.gov/document/USCG-2019-0131-0100

U.S. Department of Energy. (2022). *Offshore wind energy strategies*. https://www.energy.gov/sites/default/files/2022-01/offshore-wind-energy-strategies-report-january-2022.pdf

Vabø, R., Olsen, K., & Huse, I. (2002). The effect of vessel avoidance of wintering Norwegian spring spawning herring. *Fisheries Research*, *58*, 59–77. https://doi.org/10.1016/S0165-7836(01)00360-5

van Berkel, J., Burchard, H., Christensen, A., Mortensen, L. O., Petersen, O. S., & Thomsen, F. (2020). The effects of offshore wind farms on hydrodynamics and implications for fishes. *Oceanography*, *33*, 108–117. https://doi.org/10.5670/oceanog.2020.410

van Putten, I. E., Plagányi, ÉE, Booth, K., Cvitanovic, C., Kelly, R., Punt, A. E., & Richards, S. A. (2018). A framework for incorporating sense of place into the management of marine systems. *Ecology and Society*, *23*, Article 4. https://doi.org/10.5751/ES-10504-230404

Vilas, D., Buszowski, J., Sagarese, S., Steenbeek, J., Siders, Z., & Chagaris, D. (2023). Evaluating red tide effects on the west Florida shelf using a spatiotemporal ecosystem modeling framework. *Scientific Reports*, *13*, Article 2541. https://doi.org/10.1038/s41598-023-29327-z

Voet, H. E. E., Van Colen, C., & Vanaverbeke, J. (2022). Climate change effects on the ecophysiology and ecological functioning of an offshore wind farm artificial hard substrate community. *The Science of the Total Environment*, *810*, Article 152194. https://doi.org/10.1016/j.scitotenv.2021.152194

Voinov, A., Jenni, K., Gray, S., Kolagani, N., Glynn, P. D., Bommel, P., Prell, C., Zellner, M., Paolisso, M., Jordan, R., & Sterling, E. (2018). Tools and methods in participatory modeling: Selecting the right tool for the job. *Environmental Modelling & Software: With Environment Data News*, *109*, 232–255. https://doi.org/10.1016/j.envsoft.2018.08.028

Voinov, A., Seppelt, R., Reis, S., Nabel, J. E., & Shokravi, S. (2014). Values in socio-environmental modelling: Persuasion for action or excuse for inaction. *Environmental Modelling & Software*, *53*, 207–212. https://doi.org/10.1016/j.envsoft.2013.12.005

Wang, K., Xiao, W., He, T., & Zhang, M. (2024). Remote sensing unveils the explosive growth of global offshore wind turbines. *Renewable*

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025

and Sustainable Energy Review, 191, Article 114186. https://doi.org/10.1016/j.rser.2023.114186

Ward, C. H., & Tunnell, J. W., Jr. (2017). Habitats and biota of the Gulf of Mexico: An overview. In C. Herb Ward (Ed.), *Habitats and biota of the Gulf of Mexico: Before the Deepwater Horizon oil spill. Volume 1: Water quality, sediments, sediment contaminants, oil and gas seeps, coastal habitats, offshore plankton and benthos, and shellfish* (pp. 1–54). Springer. https://doi.org/10.1007/978-1-4939-3447-8_1

Werner, K. M., Haslob, H., Reichel, A. F., Gimpel, A., & Stelzenmüller, V. (2024). Offshore wind farm foundations as artificial reefs: The devil is in the detail. *Fisheries Research, 272*, Article 106937. https://doi.org/10.1016/j.fishres.2024.106937

White, J. W., Carr, M. H., Caselle, J. E., Washburn, L., Woodson, C. B., Palumbi, S. R., Carlson, P. M., Warner, R. R., Menge, B. A., & Barth, J. A. (2019). Connectivity, dispersal, and recruitment: connecting benthic communities and the coastal ocean. *Oceanography, 32*, 50–59. https://doi.org/10.5670/oceanog.2019.310

Wilber, D. H., Brown, L. J., Griffin, M., & Carey, D. A. (2024). American lobster *Homarus americanus* responses to construction and operation of an offshore wind farm in southern new England. *Marine Ecology Progress Series, 727*, 123–142. https://doi.org/10.3354/meps14482

Wilber, D. H., Brown, L., Griffin, M., DeCelles, G. R., & Carey, D. A. (2022a). Demersal fish and invertebrate catches relative to construction and operation of North America's first offshore wind farm. *ICES Journal of Marine Science, 79*, 1274–1288. https://doi.org/10.1093/icesjms/fsac051

Wilber, D. H., Brown, L., Griffin, M., DeCelles, G. R., & Carey, D. A. (2022b). Offshore wind farm effects on flounder and gadid dietary habits and condition on the northeastern US coast. *Marine Ecology Progress Series, 683*, 123–138. https://doi.org/10.3354/meps13957

Wilhelmsson, D., Malm, T., & Öhman, M. C. (2006). The influence of offshore windpower on demersal fish. *ICES Journal of Marine Science, 63*, 775–784. https://doi.org/10.1016/j.icesjms.2006.02.001

Williams, A. B. (1958). Substrates as a factor in shrimp distribution. *Limnology and Oceanography, 3*, 283–290. https://doi.org/10.4319/lo.1958.3.3.0283

Williams, R., Backwell, B. F., Lee, J., Patel, A., Smith, M., Hutchinson, M., Lathigara, A., Liang, W., Fang, E., Ngueyen, T., Hubbard, B., Ruas, M., Fiestas, R., Muchiri, W., Qiao, L, Vinh Bui, T., Jo, E., & Weekes, N. (2023). *Global offshore wind report 2023*. Global Wind Energy Council.

Williams, J. A., Johnson, K., & Lowther, A. (2024). *SEDAR 87: Commercial fishery landings and effort figures for white, pink, and brown shrimp in the US Gulf of Mexico, 1960–2021* (SEDAR87-DW-16). SEDAR.

Wilson, D. C., & McCay, B. J. (1998). How the participants talk about "participation" in mid-Atlantic fisheries management. *Ocean & Coastal Management, 41*, 41–61. https://doi.org/10.1016/S0964-5691(98)00080-5

Yergin, D. (1991). *The prize: The epic quest for oil, money, and power*. Simon and Schuster.

Zein-Eldin, Z. P., & Aldrich, D. V. (1965). Growth and survival of postlarval *Penaeus aztecus* under controlled conditions of temperature and salinity. *The Biological Bulletin, 129*, 199–216. https://doi.org/10.2307/1539778

Zein-Eldin, Z. P., & Renaud, M. L. (1986). Inshore environmental effects on brown shrimp, *Penaeus aztecus*, and white shrimp, *P. setiferus*, populations in coastal waters, particularly of Texas. *Marine Fisheries Review, 48*, 9–19. https://spo.nmfs.noaa.gov/sites/default/files/pdf-content/MFR/mfr483/mfr4832.pdf

Zupan, M., Rumes, B., Vanaverbeke, J., Degraer, S., & Kerckhof, F. (2023). Long-term succession on offshore wind farms and the role of species interactions. *Diversity, 15*, Article 288. https://doi.org/10.3390/d15020288

Downloaded from https://academic.oup.com/mcf/article/17/3/vtaf009/8150955 by guest on 19 June 2025